

07/18/00  
3885 U.S. PTO

07-20-00

A

CONTINUATION APPLICATION TRANSMITTAL

U.S. PTO  
09/18/00  
07/18/00

Customer No. 20350

TOWNSEND and TOWNSEND and CREW LLP  
Two Embarcadero Center, 8<sup>th</sup> Floor  
San Francisco, California 94111-3834  
(650) 326-2400

Attorney Docket No. 11823-004920US

Client Ref No. PDL0049-20

"Express Mail" Label No. EL265820084US

Date of Deposit: July 18, 2000

ASSISTANT COMMISSIONER FOR PATENTS  
BOX PATENT APPLICATION  
Washington, D.C. 20231

I hereby certify that this is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above, addressed to:

Box Patent Application  
Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

Transmitted herewith for filing under 37 CFR 1.53(b) is the  
[ X ] continuation patent application of

By: Ron Anton

Applicant: T. YUN TSO

For: BISPECIFIC ANTIBODY EFFECTIVE TO TREAT B-CELL LYMPHOMA AND CELL LINE

[ X ] This application claims priority from each of the following Application Nos./filing dates: Application No. 08/397,411 filed March 1, 1995 and Application No. 07/859,583 filed March 27, 1992, the disclosures of which are incorporated by reference.

PLEASE CANCEL CLAIMS 1-15 and 19-33.

Enclosed are:

- [ X ] 39 pages of specification
- [ X ] 8 pages of claims
- [ X ] 1 -page Abstract
- [ X ] 14 sheets of informal drawings
- [ X ] Preliminary Amendment (2 pages)
- [ X ] Sequence Listing (20 pages) and Transmittal (1 page)
- [ X ] Request to Delete Inventors (1 page)
- [ X ] Assignment of the invention to Protein Design Labs, Inc. (6 pages)
- [ X ] Unsigned Declaration & Power of Attorney (2 pages)
- [ X ] Verified statement to establish small entity status under 37 CFR 1.9 and 37 CFR 1.27 (1 page) was filed in the prior application; small entity status is still proper and desired.
- [ X ] Information Disclosure Statement under 37 CFR 1.97 (2 pages) and PTO Form 1449 (2 pages)
- [ X ] Return Postcard

**In view of the unsigned Declaration and Power of Attorney as filed with this application and pursuant to 37 CFR §1.53(f), Applicants request deferral of the filing fee until submission of the Missing Parts of Application.**

DO NOT CHARGE THE FILING FEE AT THIS TIME.

Please Charge Deposit Account No. 20-1430 With The Following:  
Any Additional Fees Associated With This Paper Or During The Pendency Of This Application.

NO Additional Copies Of This Transmittal Sheet Are Enclosed.

Respectfully submitted,  
TOWNSEND and TOWNSEND and CREW LLP

J. Liebeschuetz

Joe Liebeschuetz  
Reg. No. 37,505  
Attorneys for Applicant

Telephone:  
(650) 326-2400

Facsimile:  
(650) 326-2422

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: J. YUN TSO

Application No.: Unassigned

Filed: Herewith

For: BISPECIFIC ANTIBODY  
EFFECTIVE TO TREAT B-CELL  
LYMPHOMA AND CELL LINE

Examiner:

Art Unit:

REQUEST TO DELETE INVENTORS

Assistant Commissioner for Patents  
Box Patent Application  
Washington, D.C. 20231

Sir:

This application is a continuation of U.S. Patent Application No. 08/397,411 filed March 1, 1995. Since T. Yun Tso is the sole inventor of the claims remaining after the preliminary amendment, the names of the following inventors in the prior application should be deleted from the current application:

George Weiner;  
Roger Gingrinch; and  
Brian Link.

Respectfully submitted,



Joe Liebeschuetz  
Reg. No. 37,505

TOWNSEND and TOWNSEND and CREW LLP  
Two Embarcadero Center, 8<sup>th</sup> Floor  
San Francisco, California 94111-3834  
Tel.: 650-326-2400  
Fax: 650-326-2422  
JOL:llr  
PA 3083342 v1

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS**  
**(37 CFR 1.9(f) & 1.27(c)) - SMALL BUSINESS CONCERN**

Applicant or Patentee: GEORGE WEINER ET AL.Serial or Patent No.: 08/397,411Filed or Issued: March 1, 1995Title: BISPECIFIC ANTIBODY EFFECTIVE TO TREAT B-CELL LYMPHOMA AND CELL LINE

I hereby declare that I am

- ☐ the owner of the small business concern identified below:  
☒ an official of the small business concern empowered to act on behalf of the concern identified below:

ORIGINAL

NAME OF SMALL BUSINESS CONCERN PROTEIN DESIGN LABS, INC.ADDRESS OF SMALL BUSINESS CONCERN 2375 Garcia Avenue, Mountain View, California 94043

I hereby declare that the above-identified small business concern qualifies as a small business concern as defined in 13 CFR 121.12, and reproduced in 37 CFR 1.9(d), for purposes of paying reduced fees to the United States Patent and Trademark Office, in that the number of employees of the concern, including those of its affiliates, does not exceed 500 persons. For purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern controls or has the power to control the other, or a third party or parties controls or has the power to control both.

I hereby declare that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention, entitled BISPECIFIC ANTIBODY EFFECTIVE TO TREAT B-CELL LYMPHOMA AND CELL LINE by inventor(s) GEORGE WEINER; ROGER GINGRICH; BRIAN LINK and J. YUN TSO described in

- ☐ the specification filed herewith  
☒ application Serial No. 08/397,411, filed March 1, 1995  
☐ Patent No. \_\_\_\_\_, issued \_\_\_\_\_

If the rights held by the above identified small business concern are not exclusive, each individual, concern or organization having rights in the invention is listed below\* and no rights to the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 CFR 1.9(c) if that person made the invention, or by any concern that would not qualify as a small business concern under 37 CFR 1.9(d), or a nonprofit organization under 37 CFR 1.9(e).

\*NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

NAME Iowa Immunotherapy InvestigatorsADDRESS 2228 Cae Drive, Iowa City, Iowa 52246

☐ INDIVIDUAL ☒ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

NAME \_\_\_\_\_

ADDRESS \_\_\_\_\_

☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b)).

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

NAME OF PERSON SIGNING

TITLE OF PERSON IF OTHER THAN OWNER

ADDRESS OF PERSON SIGNING

SIGNATURE

DATE

Cary Queen (Cary Queen)Senior Vice President622 Benvenue Ave, Los Altos, CA 94024Cary Queen5/10/95

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In re application of:

J. YUN TSO

Application No.: Unassigned

Filed: July 18, 2000

For: BISPECIFIC ANTIBODY  
EFFECTIVE TO TREAT B-CELL  
LYMPHOMA AND CELL LINE

Examiner:

Art Unit:

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

Prior to examination of the above-referenced application, please enter the following amendments and remarks.

IN THE SPECIFICATION:

On page 1, delete lines 7-10.

On page 6, at line 13, after "(A)" insert -- (SEQ ID NOS. 1 and 2) --;

same page, at line 13, after "(B)" insert -- (SEQ ID NOS. 3 and 4) --;

same page, at line 20, after "(C)" insert -- (SEQ ID NO. 5) --;

same page, at line 20, after "(E)" insert -- (SEQ ID NO. 7) --;


same page, at line 25, after "(D)" insert -- (SEQ ID NO. 6) --;

same page at line 29, after "(A)" insert -- (SEQ ID NOS. 8 and 9) --;

same page at line 29, after "(B)" insert -- (SEQ ID NOS 10 and 11) --;

same page at line 36, after "(C)" insert -- (SEQ ID NO. 12) --.

Respectfully submitted,



Joe Liebeschuetz  
Reg. No. 37,505

TOWNSEND and TOWNSEND and CREW LLP  
Two Embarcadero Center, Eighth Floor  
San Francisco, California 94111-3834  
Tel.: (650) 326-2400  
Fax: (650) 326-2422  
JOL:llr  
PA 3083733 v1

PATENT APPLICATION

BISPECIFIC ANTIBODY EFFECTIVE TO TREAT B-CELL  
LYMPHOMA AND CELL LINE

Inventors:

GEORGE WEINER of Iowa City, Iowa;  
ROGER GINGRICH of Iowa City, Iowa;  
BRIAN LINK of Carlville, Iowa; and  
J. YUN TSO of Menlo Park, California

Assignees:

IOWA IMMUNOTHERAPY INVESTIGATORS  
PROTEIN DESIGN LABS, INC.

Entity:

SMALL

TOWNSEND and TOWNSEND KHOURIE and CREW  
Steuart Street Tower, 20th Floor  
One Market Plaza  
San Francisco, California 94105  
(415) 326-2400

BISPECIFIC ANTIBODY EFFECTIVE TO TREAT B-CELL  
LYMPHOMA AND CELL LINE

CROSS-REFERENCE TO RELATED APPLICATION

This application is a continuation-in-part of USSN  
07/859,583 filed 3/27/92, which is incorporated by reference  
in its entirety for all purposes.

BACKGROUND OF THE INVENTION

Administration of monoclonal antibodies (MoAb) has  
shown promise as a new treatment modality for human  
malignancy. However, destruction of malignant cells by MoAb  
does not always occur, even after successful binding of the  
antibody to the target cell. A second approach to  
immunotherapy of malignancy involves the manipulation of the  
cellular immune system. Lymphokines, such as IL-2, can be  
used to activate both NK cells and T cells isolated from the  
blood, spleen, or malignant tumors themselves. The anti-tumor  
effects of such cells have been well documented both *in vitro*  
and *in vivo*. Toxicity of therapy based on IL-2 alone can be  
severe and may well limit the clinical utility of this  
therapy.

Immunotherapy of malignancy that attempts to combine  
the specificity of antibodies with the power of activated  
lymphocytes might be more effective and less toxic. One such  
approach is the use of bispecific antibodies to redirect  
activated T cell toxicity toward tumor cells expressing the  
target antigen (Ag.)

Various forms of bispecific antibodies have been  
produced. These include BSIgG, which are IgG molecules  
comprising two distinct heavy chains and two distinct light  
chains that are secreted by so-called "hybrid hybridomas", and  
heteroantibody conjugates produced by the chemical conjugation  
of antibodies or antibody fragments of different  
specificities.

Several investigators have evaluated anti-CD3/anti-tumor bispecific antibody structures as immunotherapeutic agents. Such studies have reported *in vitro* cytotoxicity of renal cell carcinoma, melanoma, glioma, lymphoma, leukemia and cells expressing the multidrug-resistance-related glycoprotein. IL-2-activated human peripheral lymphocytes directed by certain anti-CD3/anti-tumor-specific heteroantibody conjugates have also been reported to prevent the growth of human cancer xenografts in nude mice. Studies *in vitro*, and *in vivo* in immunodeficient mice bearing human xenografts have reported that certain bispecific antibodies are capable of blocking the growth of both tumor cells bearing certain target antigens and, to some extent, bystander tumor cells that are not recognized by the therapeutic antibody.

The cell membranes of lymphocytes are uniquely constructed and determine such diverse cellular phenotypic characteristics as the suppressor, inducer, or cytolytic function of the cell, the state of activation or stage of differentiation of the cell, and whether the cell belongs to a population that is monoclonal or polyclonal. The vast majority of cellular membrane antigens thus far described on malignant lymphocytes are represented on nonmalignant lymphocytes at some stage of differentiation or activation.

From the foregoing, it is apparent that a need exists for therapeutic agents that are targeted to an antigen found predominantly or exclusively on malignant cells, and which are capable of inducing strong cytolytic activity against such cells. The present invention fulfills this and other needs.

#### SUMMARY OF THE INVENTION

The present invention is premised on the realization that a bispecific monoclonal antibody which binds to malignant B-cell lymphomas and to T cells can be formed which effectively binds only to malignant B-cells and does not bind to normal B-cells.

Further, the present invention is premised on the realization that a bispecific antibody can be formed from a



cell line obtained from peripherally diffuse large cell lymphoma to produce a monoclonal antibody that is specific only to malignant B-cells and that this monoclonal antibody can be modified to form a bispecific antibody which also binds to killer T cells or NK cells.

The present invention is further premised on the realization that a cell line formed from a fusion of cell lines which produces an IgG antibody specific to the T cells or NK cells and a cell line which produces the IgG antibody specific to B-cell malignancies in turn produce a unique bispecific antibody that effectively binds to both malignant B-cells and T cells or NK cells thereby effectuating the lysis or destruction of the malignant B-cells.

In the preferred embodiment the cell line is derived from the fusion of a cell line producing an antibody specific to the CD3 antigen of the T cell in combination with a cell line specific to a heterodimer on the cell membrane of the malignant B-cells as explained further below.

In a further aspect, the invention provides the 1D10 antibody, which is specific for the 28/32 kDa heterodimeric protein on the surface of malignant B-cells.

The invention further provides a humanized version of the 1D10 antibody. The humanized antibody comprises a humanized heavy chain and a humanized light chain. The humanized light chain comprises three complementarity determining regions (CDR1, CDR2 and CDR3) having amino acid sequences from the corresponding complementarity determining regions of the 1D10 immunoglobulin light chain, and a variable region framework from a human kappa light chain variable region framework sequence except in at least one position selected from a first group consisting of L48, L49, L69, and L70 wherein the amino acid position is occupied by the same amino acid present in the equivalent position of the 1D10 immunoglobulin light chain variable region framework. The humanized heavy chain comprising three complementarity determining regions (CDR1, CDR2 and CDR3) having amino acid sequences from the corresponding complementarity determining regions of 1D10 immunoglobulin heavy chain, and a variable

region framework from a human heavy chain variable region framework sequence except in at least one position selected from a second group consisting of H27, H29, H30, H37, H67, H71, H78 and H83, wherein the amino acid position is occupied by the same amino acid present in the equivalent position of the mouse 1D10 immunoglobulin heavy chain variable region framework. The humanized antibody specifically binds the 28/32 kDa heterodimeric protein cells with a binding affinity having a lower limit of about  $10^7 \text{ M}^{-1}$  and an upper limit of about five-times the binding affinity of the 1D10 immunoglobulin. Preferably, the humanized light chain variable region framework is from the R3.5H5G antibody. In this case, position L43 can be substituted with the amino acid present in the equivalent position of a human kappa subgroup I consensus sequence. Preferably, the humanized heavy chain is from the heavy chain region variable framework of the IC4 antibody. In this case, position H73 can be substituted by the same amino acid present in the equivalent position of a human immunoglobulin subgroup II or IV consensus sequence.

In a further aspect, the invention provides humanized antibodies specific for the CD3 antigen. The antibodies comprise humanized heavy and light chains. The humanized light chain comprises three complementarity determining regions (CDR1, CDR2 and CDR3) having amino acid sequences from the corresponding complementarity determining regions of the M291 immunoglobulin light chain, and a variable region framework from a human kappa light chain variable region framework sequence. The humanized heavy chain comprises three complementarity determining regions (CDR1, CDR2 and CDR3) having amino acid sequences from the corresponding complementarity determining regions of M291 immunoglobulin heavy chain, and a variable region framework from a human heavy chain variable region framework sequence except in at least one position selected from a second group consisting of H30, H67, H68, H70, H72 and H74 wherein the amino acid position is occupied by the same amino acid present in the equivalent position of the mouse M291 immunoglobulin heavy

chain variable region framework. The immunoglobulin specifically binds to a CD3 antigen on the surface of T cells with a binding affinity having a lower limit of about  $10^7 \text{ M}^{-1}$  and an upper limit of about five-times the binding affinity of the M291 immunoglobulin. Preferably, the humanized light chain variable region framework is from the light chain variable region framework of the HF2-1/17 antibody in subgroup I. Preferably, the humanized heavy chain region framework is from the heavy chain region variable framework of the 21/28 antibody. In this case, position H44 can be substituted with the same amino acid present in the equivalent position of a human immunoglobulin subgroup I consensus sequence.

In a further aspect, the invention provides humanized bispecific antibodies comprising a first binding fragment that specifically binds to the CD3 antigen and a second binding fragment that specifically binds to the 28/32 kDa heterodimeric antigen on the surface of the malignant B cells. The first binding fragment comprises a humanized form of the heavy chain variable region of the M291 antibody and a humanized form of the light chain variable region of the M291 antibody. The second binding fragment, which is linked to the first binding fragment, comprising: a humanized form of the heavy chain variable region from the 1D10 antibody and a humanized form of the light chain variable region from the 1D10 antibody.

Preferably, the first and second binding fragments each further comprises a segment of a constant region fused to the respective heavy chain variable regions, and the binding fragments are linked by association of the constant regions. For, example, the binding fragments can be Fab or Fab'. When both binding fragments are Fab', the bispecific antibody is a  $\text{F(ab')}_2$ . Optionally, the first and second binding fragments further comprise first and second leucine zippers fused to the respective constant regions.

The invention further provides pharmaceutical compositions comprising the antibodies described above. Also provided are methods of treating patients suffering from

malignant B-cells employing a therapeutically effective amount of bispecific antibody as described above.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph depicting lysis of malignant B-cells by the bispecific antibody of the present invention;

Figure 2 is a graph depicting lysis of Raji cells caused by different concentrations of the antibody of the present invention;

Figure 3 is a graph depicting lysis of KH cells over a period of time by the bispecific antibody of the present invention also depicting a comparative study.

Figure 4. Amino acid sequences of the light chain (A) and the heavy chain (B) variable regions of the humanized 1D10 antibody (upper lines) and mouse 1D10 antibody (lower lines), not including the signal sequences. The three CDRs in each chain are underlined. Residues in the human framework that have been replaced with mouse amino acids or consensus human amino acids are doubly underlined. Amino acid sequences of the complete light chain and the heavy chain of the humanized 1D10 are showed in (C) and (E), respectively. The  $V_L$  domain consists of residues 1-107, and the  $C_K$  108-214. The  $V_H$  domain consists of residues 1-116, the  $C_{H1}$  117-214, the hinge 215-229, the  $C_{H2}$  230-339, and the  $C_{H3}$  domain 340-446. Amino acid sequence of the Fd-Jun in the humanized F(ab'-zipper)<sub>2</sub> of 1D10 is shown in (D). The  $V_H$  domain consists of residues 1-116, the  $C_{H1}$  domain 117-214, the modified hinge 215-234, and the Fos leucine zipper 235-273.

Figure 5. Amino acid sequences of the light chain (A) and the heavy chain (B) variable regions of the humanized M291 antibody (upper lines) and mouse M291 antibody (lower lines), not including the signal sequences. The three CDRs in each chain are underlined. Residues in the human framework that have been replaced with mouse amino acids or consensus human amino acids are doubly underlined. Amino acid sequences of the complete light chain of the humanized M291 are showed in (C). The  $V_L$  domain consists of residues 1-106, and the human  $C_K$  domain 107-213. Amino acid sequence of the Fd-Fos in the humanized F(ab'zipper)<sub>2</sub> of M291 is shown in (D). The  $V_H$

domain consists of residues 1-120, the C<sub>H</sub>1 domain 121-218, the modified hinge 219-238, and the Fos leucine zipper 239-279.

Figure 6. Construction of the plasmid pHu1D10.IgG1.rG.dE used for the expression of the humanized 1D10 IgG1.

Figure 7. (A). Displacement assay to compare the relative affinity of humanized 1D10 and murine 1D10 for the antigen. Subsaturating amounts of murine 1D10-IgG2a-FITC on Raji cells were displaced by increasing amounts of murine 1D10-IgG2a or humanized 1D10-IgG1. Raji cells were resuspended in complete media at  $2.5 \times 10^6$ /ml. Dilutions of the test (humanized 1D10-IgG1) or control (murine 1D10-IgG2a) antibody were added and incubated at 4°C for 1 hour. A fixed, subsaturating amount of murine 1D10-IgG2a-FITC was added, and the cells were incubated at 4°C for 1 hour, washed, and resuspended in 1% paraformaldehyde. The cells were then analyzed using flow cytometry. Values expressed in % inhibition of fluorescence intensity compared to no competitive antibody control. (B). Scatchard plot analysis of the binding of  $^{125}\text{I}$ -labeled humanized 1D10-IgG1 to Raji cells. Scatchard analysis was made by binding dilutions of labeled antibody to  $4 \times 10^5$  Raji cells in 0.2 ml for 90 min at 0°C. The cells were washed in binding buffer (2% horse serum in PBS containing 0.1% sodium azide) and counted. Nonspecific binding was determined by inhibiting the specific binding with an excess of nonlabeled humanized 1D10-IgG1. The apparent  $K_a$  and the number of binding sites were calculated from the slope and the X axis intercept, respectively, of the Scatchard plot.

Figure 8. (A). Antibody-dependent cell-mediated cytotoxicity (ADCC) capability by various 1D10 isotypes.  $^{51}\text{Cr}$ -labeled Raji human lymphoma cells were used as targets for (▲) murine 1D10-IgG1, (●) murine 1D10-IgG2a, or (■) humanized 1D10-IgG1 and human peripheral mononuclear as effector cells. The effector:target ratio was 40:1. Spontaneous release was less than 20% of total release. Bars represent SEM. (B). Complement-mediated cytotoxicity by various 1D10 isotypes.  $^{51}\text{Cr}$ -labeled Raji human lymphoma cells were used as targets for (▲) murine 1D10-IgG1, (●) murine

1D10-IgG2a, or (■) humanized 1D10-IgG1 and human sera from a normal subject as complement. Spontaneous release was less than 20% of total release. Bars represent SEM.

Figure 9. Schematic diagrams of the plasmids pHu1D10-Jun.rG.dE and pHuM291-Fos.rG.dE for the expression of Hu1D10-Jun and HuM291-Fos F(ab'-zipper)<sub>2</sub>. The constructions of these two plasmids were similar to that of pHu1D10.IgG1 in Figure 6 except for the replacement of the C<sub>H</sub>2 and C<sub>H</sub>3 exons by the leucine zipper sequences Jun and Fos. The polyadenylation signal for the Fd-zipper transcript is from the 3' noncoding sequence of mouse IgG2a gene (see Kostelny et al., *J. Immunol.* 148, 1547 (1992)).

Figure 10. (A). The sequence of the modified human IgG1 hinge used in the hinge-zipper fusion. Two residues Lys-Cys (underlined) were inserted in the modified hinge. The first Cys in this modified hinge forms disulfide bond with the light chain, and the last three Cys residues form inter-heavy chain disulfides. For comparison, hinge sequences of the human IgG1 (B) and the mouse IgG2a (C) are also shown. All three Cys residues in the mouse IgG2a hinge are used for inter-heavy chain disulfides. After the insertion of Lys-Cys, the modified hinge and the mouse IgG2a hinge have extensive sequence homology near the COOH-terminus.

Figure 11. (A). Displacement assay to compare the relative affinity of HuM291-Fos and M291 for their antigen. Subsaturating amounts of murine M291-FITC on human T cells were displaced by increasing amounts of murine M291 or HuM291-Fos. T cells were resuspended in complete media at  $2.5 \times 10^6$ /ml. Dilutions of the test (HuM291-Fos) or control (murine M291) antibody were added and incubated at 4°C for 1 hour. A fixed, subsaturating amount of murine M291-FITC was added, and the cells were incubated at 4°C for 1 hour, washed, and resuspended in 1% paraformaldehyde. The cells were then analyzed using flow cytometry. Values expressed in % inhibition of fluorescence intensity compared to no competitive antibody control. (B). Scatchard plot analysis of the binding of <sup>125</sup>I-labeled HuM291-Fos to activated human T cells. Scatchard analysis was made by binding dilutions of

labeled antibody to  $4 \times 10^5$  T cells in 0.2 ml for 90 min at 0°C. The cells were washed in binding buffer (2% horse serum in PBS containing 0.1% sodium azide) and counted. Nonspecific binding was determined by inhibiting the specific binding with an excess of nonlabeled HuM291-Fos. The apparent  $K_a$  and the number of binding sites were calculated from the slope and the X axis intercept, respectively, of the Scatchard plot.

Figure 12. Bispecific antibody induced T cell mediated lysis of 1D10 positive cells. T cells in human PBL were activated by anti-CD3 antibody OKT3 and expanded by culturing them in IL-2. Target cells were labeled with  $^{51}\text{Cr}$  and washed. T cells and labeled target cells at effector: target ratio of 25:1 were plated in V bottom microtiter plates. Antibodies at desired concentration were added. Antibodies used were: Hu1D10-Jun, HuM291-Fos, the mouse bispecific IgG 1DT3-D, and the humanized bispecific F(ab'-zipper)<sub>2</sub> Hu1D10-Jun x HuM291-Fos. Plates were incubated at 37°C for 4 hours, centrifuged, and target cell lysis was measured by determining the amount of  $^{51}\text{Cr}$  released. Percentages of specific release in this cytotoxicity assay were calculated as: {Counts released by antibody minus counts released without added antibody}/{Counts released by 0.1% SDS minus counts released without added antibody} x 100.

#### DEFINITIONS

The term "substantial identity" or "substantial homology" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 65 percent sequence identity, preferably at least 80 or 90 percent sequence identity, more preferably at least 95 percent sequence identity or more (e.g., 99 percent sequence identity). Preferably, residue positions which are not identical differ by conservative amino acid substitutions.

For purposes of classifying amino acids substitutions as conservative or nonconservative, amino acids are grouped as follows: Group I (hydrophobic sidechains): norleucine, met, ala, val, leu, ile; Group II (neutral

hydrophilic side chains): cys, ser, thr; Group III (acidic side chains): asp, glu; Group IV (basic side chains): asn, gln, his, lys, arg; Group V (residues influencing chain orientation): gly, pro; and Group VI (aromatic side chains): trp, tyr, phe. Conservative substitutions involve substitutions between amino acids in the same class. Non-conservative substitutions constitute exchanging a member of one of these classes for a member of another.

Amino acids from the variable regions of the mature heavy and light chains of immunoglobulins are designated Hx and Lx respectively, where x is a number designating the position of an amino acids according to the scheme of Kabat, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, MD, 1987 and 1991). Kabat lists many amino acid sequences for antibodies for each subclass, and lists the most commonly occurring amino acid for each residue position in that subclass. Kabat uses a method for assigning a residue number to each amino acid in a listed sequence, and this method for assigning residue numbers has become standard in the field. Kabat's scheme is extendible to other antibodies not included in his compendium by aligning the antibody in question with one of the consensus sequences in Kabat. The use of the Kabat numbering system readily identifies amino acids at equivalent positions in different antibodies. For example, an amino acid at the L50 position of a human antibody occupies the equivalent position to an amino acid position L50 of a mouse antibody.

From N-terminal to C-terminal, both light and heavy chains comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is in accordance with the definitions of Kabat (1987) and (1991), *supra*, or Chothia & Lesk, *J. Mol. Biol.* 196:901-917 (1987); Chothia et al., *Nature* 342:878-883 (1989).

The basic antibody structural unit is known to comprise a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" chain (about 50-70 kDa). The amino-terminal portion of each chain includes a



variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function. The variable regions of each light/heavy chain pair form the antibody binding site. Thus, an intact antibody has two binding sites.

Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, and define the antibody's isotype as IgG, IgM, IgA, IgD and IgE, respectively. Within light and heavy chains, the variable and constant regions are joined by a "J" region of about 12 or more amino acids, with the heavy chain also including a "D" region of about 10 more amino acids. (See generally, *Fundamental Immunology* (Paul, W., ed., 2nd ed. Raven Press, N.Y., 1989), Ch. 7 (incorporated by reference in its entirety for all purposes).

The term epitope includes any protein determinant capable of specific binding to an immunoglobulin or T-cell receptor. Epitopic determinants usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics.

The term patient includes human and veterinary subjects.

#### DETAILED DESCRIPTION

The present invention provides bispecific antibodies, which are specific to both effector cells (T cells or natural killer cells) and to a 28/32 kDa heterodimeric antigen present on the surface of malignant B-cells. The present invention further provides hybridomas and other cells lines producing the claimed antibodies.

The 28/32 kDa antigen is found predominantly on the surface of malignant B lymphocytes and is not expressed on resting lymphocytes or B and T cells activated *in vitro* by a variety of inductive stimuli. See Gingrich et al., *Blood* 75, 2375-2387 (1990). The antigen can be expressed when lymphocytes undergo malignant transformation or, in some

cases, when they are perturbed by the Epstein-Barr virus (EBV). Normal resting and stimulated lymphocytes do not express the antigen. The antigen is also absent on hemopoietic stem cells. Although the scientific basis for the 28/32 kDa antigen being expressed predominantly or exclusively on malignant B-cells is not critical to the practice of the invention, it is believed that the antigen may represent an aberrant post-translational processing variant of the HLA-Dr antigen.

To produce the antibodies specific to malignant B-cells, a lymphoma cell line derived from a patient with peripheralizing diffuse large cell lymphoma labeled HO-85 was grown in suspension culture RPMI 1640 with 10% fetal calf serum with a doubling time of approximately 24 hours. The cell line is CD20, mu, delta (weakly), kappa, HLA Class I and II antigen positive. It does not react with monoclonal antibodies detecting CALLA, T cell, myeloid or monocytic cell antigens. The cells react with the SFR7, DR7 and B7/21 monoclonal antibodies indicating that they express DR7 and DP antigens respectively.

Female BALB/c mice, age 6-10 weeks, were given 4 to 6 intraperitoneal inoculations at two week intervals with  $5 \times 10^6$  cells from the human large cell lymphoma line as described above. The animals were killed five days after last inoculation and the spleen cells were fused with the nonsecretory murine myeloma cell line N-1. Hybridomas were selected in hypoxanthine-aminopterin-thymidine (HAT) medium after being plated in 96 well cell culture trays. After 10 days, 25 microliter aliquots were taken from each well for determination of malignant B-cell (anti-HO-85) antibody binding activity.

Malignant B-cell (anti-HO-85) antibody binding activity was determined by a whole cell, indirect radio immune assay using fresh HO-85 cells as targets. The identical assay was done using as targets RAJI (ATCC CCL86), MOLT-3 (ATCC CRL1582), HL-60 (ATCC CCL240) and fresh peripheral blood mononuclear cells. Wells that showed binding activity greater than 5 times that of tissue culture medium alone to HO-85 and

Raji but were not reactive with MOLT-3, HL-60 and peripheral blood mononuclear cells were harvested.

Cells meeting the above criteria were found to produce an antibody referred to as 1D10 and were subsequently cloned by limiting dilution. The hybridoma grows well in vitro and ascites of pristine-primed BALB/c mice.

The portion of the malignant B-cells to which 1D10 binds is a heterodimeric polypeptide which contains two proteins with a molecular weight of the alpha and beta chains being 32 kDa and 28 kDa respectively. The proteins can be obtained by solubilizing malignant B-cells such as Raji cells with detergent. Molecular weight determination is made by using iodinated cells and single dimension SDS-PAGE analysis of the MoAb6-antigen precipitate. The formation of the 1D10 antibody is discussed by Gingrich et al., *Blood* 75, 2375-2387 (1990). Other antibodies having the same or similar binding specificity to 1D10 are screened by competition binding with 1D10 to the 28/32 kDa heterodimeric antigen. Numerous types of competitive binding assays are known, for example: solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay (EIA), sandwich competition assay (see Stahli et al., *Methods in Enzymology* 9, 242-253 (1983)); solid phase direct biotin-avidin EIA (see Kirkland et al., *J. Immunol.* 137, 3614-3619 (1986)); solid phase direct labeled assay, solid phase direct labeled sandwich assay (see Harlow & Lane, "Antibodies, A Laboratory Manual," Cold Spring Harbor Press (1988)); solid phase direct label RIA using I-125 label (see Morel et al., *Molec. Immunol.* 25, 7-15 (1988)); solid phase direct biotin-avidin EIA (Cheung et al., *Virology* 176, 546-552 (1990)); and direct labeled RIA (Moldenhauer et al., *Scand. J. Immunol.* 32, 77-82 (1990)). Typically, such an assay involves the use of cells bearing the 28/32 kDa antigen, an unlabelled test immunoglobulin and a labelled reference immunoglobulin (1D10). Competitive inhibition is measured by determining the amount of label bound to the cells in the presence of the test immunoglobulin. Usually the test immunoglobulin is present in excess. Antibodies identified by competition assay (competing

antibodies) include antibodies binding to the same epitope as the reference antibody and antibodies binding to an adjacent epitope sufficiently proximal to the epitope bound by the reference antibody for steric hindrance to occur.

5           The second component for the bispecific antibodies of the invention is an antibody having specificity for an antigen on the surface of T-cells or NK cells. Human T-cell antigens likely to be suitable include CD3, CD2, CD28, CD44, C69, A13 and G1. Suitable antigens on natural killer cells  
10 include FC Gamma receptors (3G8, B73.1, LEUL1, VEP13, and AT10). Human T-cell antigens that are probably unsuitable include MHC Class I, CD4, CD8, CD18 and CD71.

Cell lines producing IgG specific to the effector cell antigens described above are commercially available or  
15 can be produced de novo (see Example 3). The OKT3 cell (ATCC CRL 8001) is a suitable source of antibodies for the CD3 antigen. Other antibodies to the CD3 antigen include WT31, WT32, anti-leu-4, UCHT-1, SPV-3TA and SPV-T3B. The CD3 site is preferred because of its presence in all T cells.

20           The antibodies of the present invention can be produced by a cell line formed by the fusion of a first component cell line producing antibodies specific for the 28/32 kDa heterodimeric antigen with a second cell line which produces an antibody specific for either T cells or natural  
25 killer cells. For example, the hybridoma producing 1D10 was fused with OKT3 as follows.

The OKT3 hybridoma cell line was selected by growing OKT3 cells sequentially in media containing 0.13 mM  
8-azaguanine, then 1.0 mM ouabain. Hybrid-hybridomas were  
30 produced by fusion (using 38% polyethylene glycol) of  $10^6$  HAT resistant, ouabain sensitive, 1D10-secreting hybridomas with  $10^6$  HAT sensitive, ouabain resistant OKT3-secreting hybridomas.

35           Fused cells were plated in HAT-ouabain media to select for hybrid-hybridomas. The HAT in this media prevented the growth of unfused OKT3 cells and the ouabain prevented the growth of unfused 1D10 cells. Thus, only hybrid-hybridomas containing genetic material from both parental hybridomas

survived. Twelve hybrid-hybridomas were isolated using this technique.

Cell lines secreting bispecific antibodies can be identified by a three-step screening procedure. For example, in analysis of hybridomas formed from fusion of 1D10 and OKT3, an initial screen was performed in which hybrid-hybridoma supernatant was added to ELISA plates coated with goat anti-mouse IgG1 antibody. After washing, alkaline phosphatase labeled goat anti-mouse IgG2a was added. Reactivity indicated the hybrid-hybridoma supernatant contained single antibody molecules with both IgG1 and IgG2a heavy chains.

An indirect immunofluorescent assay was used as a second screen for all samples that were positive on ELISA. In this second screen, hybrid-hybridoma supernatant was added separately to HO-85 (1D10 reactive) and Jurkat (OKT3 reactive) cells. Goat anti-mouse IgG-FITC was added after washing to detect the presence of bound antibody. All twelve hybrid-hybridomas secreted antibody which was capable of binding to both HO-85 and Jurkat cells. One of these hybrid-hybridomas was selected for further study. It was subcloned by limiting dilution x2, and designated 1DT3-D. This cell line was deposited on March 24, 1992 under the Budapest Treat at the American Type Culture collection, 12301 Parklawn Drive, Rockville, MD 20852 and assigned the number ATCC HB 10993.

1DT3-D was cultured *in vitro* in HB 101 media supplemented with 100  $\mu$ g L-glutamine and 100 U/ml penicillin-streptomycin. These cells were transferred to a Mini Flo-path Bioreactor hollow fiber apparatus. Antibody obtained from spent media was fractionated by HPLC cation exchange using a gradient of 0.18 to 0.5 M NaCl. The peak containing bispecific reactivity, as demonstrated by the above assays, was isolated, dialyzed against phosphate buffered saline, concentrated and used in further studies.

The bispecific antibody formed by fusion of 1D10 and OKT3 is a mouse derived monoclonal. Humanized versions of this antibody and other bispecific antibodies of the invention can also be employed as discussed in more detail below.

### Humanized Antibodies

The invention further provides humanized immunoglobulins (or antibodies). Some humanized antibodies are specific for the T-cell antigen CD3. Other humanized antibodies are specific for the 28/32 kDa heterodimer on malignant B-cells. These humanized antibodies are useful as therapeutic and diagnostic reagents in their own right or can be combined to form a humanized bispecific antibody possessing both of the binding specificities of its components. The humanized forms of immunoglobulins have variable framework region(s) substantially from a human immunoglobulin (termed an acceptor immunoglobulin) and complementarity determining regions substantially from a mouse immunoglobulin (referred to as the donor immunoglobulin). The constant region(s), if present, are also substantially from a human immunoglobulin. The humanized antibodies exhibit a specific binding affinity for their respective antigens of at least  $10^7$ ,  $10^8$ ,  $10^9$ , or  $10^{10} \text{ M}^{-1}$ . Often the upper and lower limits of binding affinity of the humanized antibodies are within a factor of three or five or ten of that of the mouse antibody from which they were derived.

#### 1. Mouse Antibodies for Humanization

The starting material for production of humanized antibody specific for the 28/32 kDa heterodimer is preferably the 1D10 mouse antibody, although other mouse antibodies, which compete with 1D10 for binding to the 28/32 kDa heterodimer can also be used. A suitable starting material for production of humanized antibody specific for CD3 is the M291 antibody whose isolation is described in Example 3.

#### (2) Selection of Human Antibodies to Supply Framework Residues

The substitution of mouse CDRs into a human variable domain framework is most likely to result in retention of their correct spatial orientation if the human variable domain framework adopts the same or similar conformation to the mouse variable framework from which the CDRs originated. This is achieved by obtaining the human variable domains from human

antibodies whose framework sequences exhibit a high degree of sequence identity with the murine variable framework domains from which the CDRs were derived. The heavy and light chain variable framework regions can be derived from the same or different human antibody sequences. The human antibody sequences can be the sequences of naturally occurring human antibodies or can be consensus sequences of several human antibodies.

Suitable human antibody sequences are identified by computer comparisons of the amino acid sequences of the mouse variable regions with the sequences of known human antibodies. The comparison is performed separately for heavy and light chains but the principles are similar for each.

### (3) Computer Modelling

The unnatural juxtaposition of murine CDR regions with human variable framework region can result in unnatural conformational restraints, which, unless corrected by substitution of certain amino acid residues, lead to loss of binding affinity. The selection of amino acid residues for substitution is determined, in part, by computer modelling. Computer hardware and software for producing three-dimensional images of immunoglobulin molecules are widely available. In general, molecular models are produced starting from solved structures for immunoglobulin chains or domains thereof. The chains to be modelled are compared for amino acid sequence similarity with chains or domains of solved three dimensional structures, and the chains or domains showing the greatest sequence similarity is/are selected as starting points for construction of the molecular model. The solved starting structures are modified to allow for differences between the actual amino acids in the immunoglobulin chains or domains being modelled, and those in the starting structure. The modified structures are then assembled into a composite immunoglobulin. Finally, the model is refined by energy minimization and by verifying that all atoms are within appropriate distances from one another and that bond lengths and angles are within chemically acceptable limits.

(4) Substitution of Amino Acid Residues

As noted *supra*, the humanized antibodies of the invention comprise variable framework region(s) substantially from a human immunoglobulin and complementarity determining regions substantially from a mouse immunoglobulin (e.g., 1D10 or M291). Having identified the complementarity determining regions of mouse antibodies and appropriate human acceptor immunoglobulins, the next step is to determine which, if any, residues from these components should be substituted to optimize the properties of the resulting humanized antibody. In general, substitution of human amino acid residues with murine should be minimized, because introduction of murine residues increases the risk of the antibody eliciting a HAMA response in humans. Amino acids are selected for substitution based on their possible influence on CDR conformation and/or binding to antigen. Investigation of such possible influences is by modelling, examination of the characteristics of the amino acids at particular locations, or empirical observation of the effects of substitution or mutagenesis of particular amino acids.

When an amino acid differs between a mouse variable framework region and an equivalent human variable framework region, the human framework amino acid should usually be substituted by the equivalent mouse amino acid if it is reasonably expected that the amino acid:

- (1) noncovalently contacts antigen directly, or
- (2) is adjacent to a CDR region or otherwise interacts with a CDR region (e.g., is within about 4-6 Å of a CDR region).

Other candidates for substitution are acceptor human framework amino acids that are unusual for a human immunoglobulin at that position. These amino acids can be substituted with amino acids from the equivalent position of more typical human immunoglobulins. Alternatively, amino acids from equivalent positions in the mouse antibody can be introduced into the human framework regions when such amino acids are typical of human immunoglobulin at the equivalent positions.



In general, substitution of all or most of the amino acids fulfilling the above criteria is desirable.

Occasionally, however, there is some ambiguity about whether a particular amino acid meets the above criteria, and

alternative variant immunoglobulins are produced, one of which has that particular substitution, the other of which does not.

The humanized antibodies of the invention that are derived from the mouse 1D10 antibody usually contain a substitution of a human kappa light chain framework residue with a corresponding mu MAb 1D10 residue in at least 1, 2, 3 or 4 of the following positions: L48, L49, L69 and L70. The humanized antibodies also usually contain a substitution of a human heavy chain framework residue in at least 1, 2, 3, 4, 5, 6, 7, or 8 of the following positions H27, H29, H30, H37, H67, H71, H78 and H83. In preferred embodiments when the human light chain acceptor immunoglobulin is R3.5HG, the light chain also contains a substitution at position 43. This position is substituted with the amino acid from the equivalent position of a human immunoglobulin having a more typical amino acid residues or from a consensus sequence of such human immunoglobulins. Similarly, when the human heavy chain acceptor immunoglobulin is IC4, the heavy chain also contains a substitution at position 73.

The humanized antibodies of the invention that are derived from mouse M291 antibody contain no substitution of a human kappa light chain framework residue if the light chain acceptor is HF2-1/17. The humanized antibodies also usually contain a substitution of a human heavy chain framework in at least 1, 2, 3, 4, 5 and 6 of the following positions: H30, H67, H68, H70, H72 and H74. In preferred embodiments, when the heavy chain acceptor immunoglobulin is 21/28, the light chain also contains a substitution at position 44. This position is substituted with the amino acid from the equivalent position of a human immunoglobulin having a more typical amino acid residue or from a consensus sequence of such human immunoglobulin.

Usually the CDR regions in humanized antibodies are substantially identical, and more usually, identical to the

corresponding CDR regions in the mouse antibody from which they were derived. Although not usually desirable, it is sometimes possible to make one or more conservative amino acid substitutions of CDR residues without appreciably affecting the binding affinity of the resulting humanized immunoglobulin. Occasionally, substitutions of CDR regions can enhance binding affinity.

Other than for the specific amino acid substitutions discussed above, the framework regions of humanized immunoglobulins are usually substantially identical, and more usually, identical to the framework regions of the human antibodies from which they were derived. Of course, many of the amino acids in the framework region make little or no direct contribution to the specificity or affinity of an antibody. Thus, many individual conservative substitutions of framework residues can be tolerated without appreciable change of the specificity or affinity of the resulting humanized immunoglobulin. However, in general, such substitutions are undesirable.

#### (5) Production of Variable Regions

Having conceptually selected the CDR and framework components of humanized immunoglobulins, a variety of methods are available for producing such immunoglobulins. Because of the degeneracy of the code, a variety of nucleic acid sequences will encode each immunoglobulin amino acid sequence. The desired nucleic acid sequences can be produced by *de novo* solid-phase DNA synthesis or by PCR mutagenesis of an earlier prepared variant of the desired polynucleotide. All nucleic acids encoding the antibodies described in this application are expressly included in the invention.

#### (6) Selection of Constant Region

The variable segments of humanized antibodies produced as described *supra* are typically linked to at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. Human constant region DNA sequences can be isolated in accordance with well-known

procedures from a variety of human cells, but preferably immortalized B-cells (see Kabat et al., *supra*, and WO87/02671). Ordinarily, the antibody will contain both light chain and heavy chain constant regions. The heavy chain constant region usually includes CH1, hinge, CH2, CH3, and, sometimes, CH4 regions.

The humanized antibodies include antibodies having all types of constant regions, including IgM, IgG, IgD, IgA and IgE, and any isotype, including IgG1, IgG2, IgG3 and IgG4. When it is desired that the humanized antibody exhibit cytotoxic activity, the constant domain is usually a complement-fixing constant domain and the class is typically IgG<sub>1</sub>. When such cytotoxic activity is not desirable, the constant domain may be of the IgG<sub>2</sub> class. The humanized antibody may comprise sequences from more than one class or isotype.

#### (7) Expression Systems

Nucleic acids encoding humanized light and heavy chain variable regions, optionally linked to constant regions, are inserted into expression vectors. The light and heavy chains can be cloned in the same or different expression vectors. The DNA segments encoding immunoglobulin chains are operably linked to control sequences in the expression vector(s) that ensure the expression of immunoglobulin polypeptides. Such control sequences include a signal sequence, a promoter, an enhancer, and a transcription termination sequence (see Queen et al., *Proc. Natl. Acad. Sci. USA* 86, 10029 (1989); WO 90/07861; Co et al., *J. Immunol.* 148, 1149 (1992), which are incorporated herein by reference in their entirety for all purposes).

#### C. Fragments of Humanized Antibodies

The humanized antibodies of the invention include fragments as well as intact antibodies. Typically, these fragments compete with the intact antibody from which they were derived for antigen binding. The fragments typically bind with an affinity of at least  $10^7 \text{ M}^{-1}$ , and more typically

10<sup>8</sup> or 10<sup>9</sup> M<sup>-1</sup> (i.e., within the same ranges as the intact antibody). Humanized antibody fragments include separate heavy chains, light chains Fab, Fab' F(ab')<sub>2</sub>, and Fv. Fragments are produced by recombinant DNA techniques, or by  
 5 enzymic or chemical separation of intact immunoglobulins.

#### Recombinant Bispecific Antibodies

The methods discussed above for forming bispecific antibodies from antibodies produced by hybridoma cells can  
 10 also be applied or adapted to production of bispecific antibodies from recombinantly expressed antibodies such as the humanized versions of 1D10 and M291. For example, bispecific antibodies can be produced by fusion of two cell lines respectively expressing the component antibodies.

15 Alternatively, the component antibodies can be co-expressed in the same cell line. Bispecific antibodies can also be formed by chemical crosslinking of component recombinant antibodies.

Component recombinant antibodies can also be linked genetically. In one approach, a bispecific antibody is  
 20 expressed as a single fusion protein comprising the four different variable domains from the two component antibodies separated by spacers. For example, such a protein might comprise from one terminus to the other, the VL region of the first component antibody, a spacer, the VH domain of the first  
 25 component antibody, a second spacer, the VH domain of the second component antibody, a third spacer, and the VL domain of the second component antibody. See, e.g., Segal et al., *Biologic Therapy of Cancer Updates* 2, 1-12 (1992).

In a further approach, bispecific antibodies are  
 30 formed by linking component antibodies to leucine zipper peptides. See generally copending application 11823-003200 (07/801,798, filed November 29, 1991; Kostelny et al., *J. Immunol.* 148, 1547-1553 (1992) (incorporated by reference in their entirety for all purposes). Leucine zippers have the  
 35 general structural formula (Leucine-X<sub>1</sub>-X<sub>2</sub>-X<sub>3</sub>-X<sub>4</sub>-X<sub>5</sub>-X<sub>6</sub>)<sub>n</sub>, where X may be any of the conventional 20 amino acids (*Proteins, Structures and Molecular Principles*, (1984) Creighton (ed.), W.H. Freeman and Company, New York), but are most likely to be

amino acids with high  $\alpha$ -helix forming potential, for example, alanine, valine, aspartic acid, glutamic acid, and lysine (Richardson and Richardson, *Science* 240, 1648 (1988)), and n may be 3 or greater, although typically n is 4 or 5. The leucine zipper occurs in a variety of eukaryotic DNA-binding proteins, such as GCN4, C/EBP, c-fos gene product (Fos), c-jun gene product (Jun), and c-myc gene product. In these proteins, the leucine zipper creates a dimerization interface wherein proteins containing leucine zippers may form stable homodimers and/or heterodimers.

The leucine zippers for use in the present invention preferably have pairwise affinity. Pairwise affinity is defined as the capacity for one species of leucine zipper, for example, the Fos leucine zipper, to predominantly form heterodimers with another species of leucine zipper, for example, the Jun leucine zipper, such that heterodimer formation is preferred over homodimer formation when two species of leucine zipper are present in sufficient concentrations. See Schuemann et al., *Nucleic Acids Res.* 19, 739 (1991). Thus, predominant formation of heterodimers leads to a dimer population that is typically 50 to 75 percent, preferentially 75 to 85 percent, and most preferably more than 85 percent heterodimers. When amino-termini of the synthetic peptides each include a cysteine residue to permit intermolecular disulfide bonding, heterodimer formation occurs to the substantial exclusion of homodimerization.

In the formation of bispecific antibodies, binding fragments of the component antibodies are fused in-frame to first and second leucine zippers. Suitable binding fragments including Fv, Fab, Fab', or the heavy chain. The zippers can be linked to the heavy or light chain of the antibody binding fragment and are usually linked to the C-terminal end. If a constant region or a portion of a constant region is present, the leucine zipper is preferably linked to the constant region or portion thereof. For example, in a Fab'-leucine zipper fusion, the zipper is usually fused to the C-terminal end of the hinge. The inclusion of leucine zippers fused to the respective component antibody fragments promotes formation of

heterodimeric fragments by annealing of the zippers. When the component antibodies include portions of constant regions (e.g., Fab' fragments), the annealing of zippers also serves to bring the constant regions into proximity, thereby promoting bonding of constant regions (e.g., in a F(ab')<sub>2</sub> fragment). Typical human constant regions bond by the formation of two disulfide bonds between hinge regions of the respective chains. This bonding can be strengthened by engineering additional cysteine residue(s) into the respective hinge regions allowing formation of additional disulfide bonds.

Leucine zippers linked to antibody binding fragments can be produced in various ways. For example, polynucleotide sequences encoding a fusion protein comprising a leucine zipper can be expressed by a cellular host or *in vitro* translation system. Alternatively, leucine zippers and/or antibody binding fragments can be produced separately, either by chemical peptide synthesis, by expression of polynucleotide sequences encoding the desired polypeptides, or by cleavage from other proteins containing leucine zippers, antibodies, or macromolecular species, and subsequent purification. Such purified polypeptides can be linked by peptide bonds, with or without intervening spacer amino acid sequences, or by non-peptide covalent bonds, with or without intervening spacer molecules, the spacer molecules being either amino acids or other non-amino acid chemical structures. Regardless of the method or type of linkage, such linkage can be reversible. For example, a chemically labile bond, either peptidyl or otherwise, can be cleaved spontaneously or upon treatment with heat, electromagnetic radiation, proteases, or chemical agents. Two examples of such reversible linkage are: (1) a linkage comprising a Asn-Gly peptide bond which can be cleaved by hydroxylamine, and (2) a disulfide bond linkage which can be cleaved by reducing agents.

Component antibody fragment-leucine zippers fusion proteins can be annealed by co-expressing both fusion proteins in the same cell line. Alternatively, the fusion proteins can be expressed in separate cell lines and mixed *in vitro*. If

the component antibody fragments include portions of a constant region (e.g., Fab' fragments), the leucine zippers can be cleaved after annealing has occurred. The component antibodies remain linked in the bispecific antibody via the constant regions.

### Therapeutic Methods

Pharmaceutical compositions comprising bispecific antibodies of the present invention are useful for parenteral administration, i.e., subcutaneously, intramuscularly and particularly, intravenously. The compositions for parenteral administration commonly comprise a solution of the antibody or a cocktail thereof dissolved in an acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers can be used, e.g., water, buffered water, 0.4% saline, 0.3% glycine and the like. These solutions are sterile and generally free of particulate matter. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, toxicity adjusting agents and the like, for example sodium acetate, sodium chloride, potassium chloride, calcium chloride, sodium lactate. The concentration of the bispecific antibodies in these formulations can vary widely, i.e., from less than about 0.01%, usually at least about 0.1% to as much as 5% by weight and will be selected primarily based on fluid volumes, and viscosities in accordance with the particular mode of administration selected.

A typical composition for intravenous infusion can be made up to contain 250 ml of sterile Ringer's solution, and 10 mg of bispecific antibody. See *Remington's Pharmaceutical Science* (15th Ed., Mack Publishing Company, Easton, Pennsylvania, 1980).

The compositions containing the present bispecific antibodies or a cocktail thereof can be administered for prophylactic and/or therapeutic treatments. In therapeutic application, compositions are administered to a patient already affected by malignant B-cells (e.g., acute

lymphoblastic leukemia, B-cell lymphoma, chronic lymphocytic leukemia and multiple myeloma) in an amount sufficient to cure or at least partially arrest the condition and its complications. An amount adequate to accomplish this is defined as a "therapeutically effective dose." Amounts effective for this use will depend upon the severity of the condition and the general state of the patient's own immune system, but generally range from about 0.01 to about 100 mg of bispecific antibody per dose, with dosages of from 0.1 to 50 mg and 1 to 10 mg per patient being more commonly used. Single or multiple administrations on a daily, weekly or monthly schedule can be carried out with dose levels and pattern being selected by the treating physician.

In prophylactic applications, compositions containing the bispecific antibodies or a cocktail thereof are administered to a patient who is at risk of developing the disease state to enhance the patient's resistance. Such an amount is defined to be a "prophylactically effective dose." In this use, the precise amounts again depend upon the patient's state of health and general level of immunity, but generally range from 0.1 to 100 mg per dose, especially 1 to 10 mg per patient.

In some methods of treatment, bispecific antibodies are administered with a second agent (e.g., an interleukin) in an amount sufficient to active effector cells thereby augmenting their cytotoxicity to malignant B-cells compared with the administration of bispecific antibody alone. Interleukin-2 at a dosage of about 500,000 U/kg is suitable. Combination therapy is particularly appropriate when the bispecific antibody being administered is a F(ab')<sub>2</sub> fragment.

The monospecific 1D10 antibody (particularly the humanized form) is also suitable for therapeutic administration to patients suffering from, or at risk of, B-cell malignancies. Optionally, the antibody is conjugated to a radiolabel or toxin. The monospecific M291 antibody (particularly the humanized form) can be used as an immunosuppressant in treatment of diseases and disorders of the immune system such as host vs. graft disease, graft vs.



host disease, autoimmune diseases, and inflammation. See, e.g., Cosimi et al., *N. Engl. J. Med.* 305, 308 (1981); Russel et al., *Annu. Rev. Med.* 35, 63 (1984). The dosages and pharmaceutical excipients for administration of monospecific antibodies are similar to those for the bispecific antibodies.

#### Diagnostic Methods

The M291 and 1D10 antibodies (both mouse and humanized forms) are also useful in diagnostic methods. The 1D10 antibody (and other antibodies binding to the same or similar epitope) is useful for diagnosing the presence of malignant B cells and monitoring the efficacy of treatments thereto. The antibody is also useful for research purposes to identify and type cells of certain lineages and developmental origins. The M291 antibody is useful for diagnostic purposes in immunologically monitoring of patients (see, e.g., Cosimi et al., *supra*) and for research purposes in classifying leukocyte subtypes, e.g., as part of an antibody panel. Methods of diagnosis can be performed *in vitro* using a cellular sample (e.g., blood sample, lymph node biopsy or tissue) from a patient or can be performed by *in vivo* imaging.

#### EXAMPLE 1

The ability of 1DT3-D to induce the elimination of malignant B cells by T cells was evaluated *in vitro*. The assay used was a  $^{51}\text{Cr}$ -chromium-release cytotoxicity assay. Target malignant B cells ( $10^7$  cells in 1 ml) were labeled during a 1 hour incubation with  $100\mu\text{Ci } ^{51}\text{Cr}$ . T cells from normal donors were incubated *in vitro* with IL-2 or IL-2 and anti CD3 antibody for 3-7 days before use as effector cells. T cells were added to  $^{51}\text{Cr}$ -labeled malignant B cells along with antibody. This mixture was incubated for 4 hours, and cell free supernatant was removed and evaluated for the presence of released  $^{51}\text{Cr}$  by gamma counting. Maximum release was determined by evaluating supernatant obtained from wells that had been treated with detergent (NP-40) that induces the lysis of all cells. Background release was determined by evaluating  $^{51}\text{Cr}$  levels from samples that had target malignant

B cells and T cells but no antibody. Specific release of  $^{51}\text{Cr}$  indicated lysis of the  $^{51}\text{Cr}$ -containing target cells, and was calculated using the following formula.

$$\frac{\text{Sample Release} - \text{Background Release}}{\text{Maximum Release} - \text{Background Release}} \times 100$$

Fig. 1 shows 1DT3-D induced the lysis of a large number of different malignant B cells including Raji (a cell line established from a patient with Burkitt's lymphoma), HO-85 (a large cell lymphoma cell line), 697 (a pre-B acute lymphoblastic leukemia cell line) and KH (fresh lymphocytes obtained from a patient with chronic lymphocytic leukemia). The T-cell target cell ratio was 10:1 and the antibody concentration was  $5\mu\text{g/ml}$ . Target lysis was not seen when monospecific antibody was used.

Fig. 2 shows 1DT3-D can induce significant lysis of raji cells at low T cell: Raji cell ratios (less than 1:1) and at low antibody concentrations (less than  $0.1\text{ ug/ml}$ ). Similar results were seen with other target cell lines.

Fig. 3 shows 1DT3-D-induced T-cell-mediated lysis of fresh KH cells was noted after long incubation times.

The bispecific antibody of the present invention can also be produced simply by taking the Fab or  $\text{F(ab')}_2$  fragments of the 1D10 antibody fusing these with portions of the OKT3 antibody to form a bispecific antibody of the present invention. Alternatively, bispecific antibodies, recognizing 1D10 and an antigen on natural killer cells or T cells, can be produced, by synthetic or genetic engineering techniques.

A benefit of the claimed bispecific antibodies is their ability to recognize malignant B-cells and distinguish these from non-malignant B-cells. Thus, therapy using the bispecific antibodies of the present invention is significantly less damaging than therapy using, for example, a non-specific antibody such as anti-CD19 antibody B4.

Further, as shown by the data described in the example, the antibody of the present invention induces significant lysis of malignant B-cells at relatively low T cell ratios. Fig. 2 shows that malignant to T cell ratios of less than 1:1 with relatively low antibody concentrations

of less than 0.1 micrograms per ml provides significant destruction of the malignant cells. This is particularly important since it reduces the dependency on the concentration of T cells available in the patient. Further, it also reduces the amount of antibody required, thereby limiting any potential side effects.

Example 2: In Vivo Efficacy of IDT3-D Bispecific Antibody

This example describes an *in vivo* trial of the bispecific antibody 1DT3-D. Normal donor human peripheral blood lymphocytes were activated *in vitro* in the presence of OKT3 (2  $\mu\text{g/ml}$ ), and recombinant IL-2 (300  $\mu\text{g/ml}$ ). CB-17 *scid/scid* mice (Itoh et al., *Cancer* 72, 2686-2694 (1993)) were injected subcutaneously with  $5 \times 10^6$  Raji cells mixed with  $5 \times 10^6$  activated lymphocytes. 24-hr later, mice were injected with bispecific antibody, a monospecific antibody component of the bispecific antibody or no antibody. Mice were examined daily for the development of tumors of at least 0.5 cm at the site of tumor injection. Mice remaining tumor-free after 60 days were scored as negative and mice developing tumors within 60 days as positive. Control untreated mice always developed tumors within 21-28 days.

In a first experiment, 5 mice were treated with 10  $\mu\text{g/mouse}$  of bispecific antibody 24 hours after inoculation with the mixture of malignant cells and activated human T-cells. A control group of 5 mice was inoculated with vehicle only. The tumor occurrence (*i.e.*, development of a tumor of at least 0.5 cm within at least eight weeks) in the treated and control groups was as follows:

<u>Group</u>	<u>No tumors</u>	<u>Tumors</u>	<u>Total</u>
Treatment	4	1	5
Control	0	5	5

Using the Fischer's one-sided exact test, bispecific antibody treatment prolonged disease-free survival with a p value of 0.024.

A second experiment was designed to compare the anti-tumor effects of bispecific antibody with monospecific anti-CD3 and monospecific 1D10 at a dose of 10  $\mu\text{g/mouse}$  in

mice inoculated with tumor and T-cells as outlined above. Group 2 mice received monospecific 1D10 and monospecific OKT3, Group 3 received bispecific antibody and the control group received vehicle only. Group 4 mice also received bispecific antibody at a concentration of 10  $\mu\text{g}/\text{mouse}$ , but these mice had previously been inoculated with unactivated T-cells as distinct from all other groups which received activated T-cells.

Group	No tumors	Tumors	Total
Control	0	5	5
2	5	0	5
3	5	0	5
4	2	3	5

Fisher's exact test for general two-way tables (Agresti, *Categorical Data Analysis* (Wiley, NY, 1990), pp. 64-65) was used to test the null hypothesis that the occurrence rates in the four groups are equal. There is a highly significant difference among the groups ( $p=0.001$ ). Pairwise exact tests comparing the control group to each of groups 2, 3, and 4 were also carried out. The corresponding one-sided  $p$ -values are 0.004, 0.004, and 0.222. Thus, groups 2 and 3 are both significantly different from the control group. It was concluded that at a dose of 10  $\mu\text{g}/\text{mouse}$  treatment with bispecific antibody or a combination of both component monospecific antibodies prolonged tumor-free survival.

In a third experiment, a dose-response study was performed to test the anti-tumor effects of varying dosages of bispecific antibody. Separate groups of mice were respectively treated with dosages of 0.4, 2 or 10  $\mu\text{g}/\text{mouse}$  bispecific antibody or vehicle.

Group	No tumors	Tumors	Total
Control	0	5	5
0.4	1	4	5
2	5	0	5
10	4	1	5

The Cochran-Armitage trend test, (Agresti, *Categorical Data Analysis* (Wiley, NY, 1990), pp. 100-102, 118-119) was used to

test the null hypothesis that the occurrence rates in the four groups are equal, versus the alternative hypotheses of a linear trend. Using equally-spaced scores, the p-value is 0.001; using the scores 0, 0.4, 2, and 10, the p-value is 0.0164. Both sets of scores indicate a significant trend in the proportions. These results show that the bispecific antibody is effective to prolong survival time and that mice receiving larger doses (10  $\mu$ g and 2  $\mu$ g) have improved tumor-free survival.

A fourth experiment was designed to compare monospecific OKT3 and 1D10 to bispecific antibody at a dose of 2  $\mu$ g antibody/mouse.

<u>Group</u>	<u>No tumors</u>	<u>Tumors</u>	<u>Total</u>
Control	0	5	5
2	0	5	5
3	1	4	5
4	4	1	5

Tumor-free survival of mice treated with monospecific OKT3 (Group 2) and monospecific 1D10 (Group 3) was not significantly different from control, whereas mice treated with bispecific antibody (Group 4) had prolonged survival using the Fisher's exact test for general two-way tables.

These data indicate that systemic administration of 1DT3-D kills and/or prevents the development of malignant B-cells *in vivo* and that a dose of 2  $\mu$ g/animal, bispecific antibody therapy is more effective than monospecific antibody therapy.

### Example 3: Generation of a monoclonal antibody against the human CD3 antigen

The 1DT3-D antibody described in Example 1 incorporated OKT3 as the binding moiety having an affinity for effector cells. The present example describes the isolation of an alternative antibody, M291, for use as the effector-cell binding component in a bispecific antibody.

Human peripheral blood mononuclear cells (PBMC) were activated with PHA and IL-2 to expand T cells. Activated T

cells were used as immunogens in Balb/C mice. Hybridomas were generated from the spleens of these mice by standard methods. These hybridomas were screened for antibodies that could stimulate PBMC to proliferate *in vitro*. Anti-CD3 antibodies with the appropriate Fc cause T cells in PBMC to proliferate. One of these hybridomas, M291, was isolated and found to secrete an antibody of the isotype IgG2a/kappa that could activate T cells to proliferate. The purified antibody M291 competes with another anti-human CD3 antibody, OKT3, (IgG2a/kappa) for binding to human T cells, showing that the epitopes recognized by the respective antibodies are closely spaced. M291 is thus an antibody having the specificity against the human CD3 complex.

#### Example 4: Humanization of 1D10 and M291 Antibodies

This example describes the separate humanization procedures for the 1D10 and M291 antibodies.

##### (1) Cloning of 1D10 and M291 V region cDNAs

Heavy and light V domain cDNAs for 1D10 and M291 were cloned using an anchored PCR method (see Loh et al., *Science* 243, 217 (1989)). cDNAs were first synthesized by reverse transcriptase after priming polyA<sup>+</sup> RNAs from the hybridoma cells with oligo dT. A tail of dGs was added to the 3' terminus of the cDNA by terminal deoxynucleotidyl transferase. The V domains were then amplified by PCR with 3' primers that hybridized to the C regions and 5' primers that hybridized to the G-tails. Several independent H and L chain clones were sequenced to ensure no sequence mistakes were introduced by PCR. For 1D10, the V domains were expressed as an antibody of the mouse isotype IgG2a/kappa by transfecting the genes in suitable vectors into the myeloma cell line SP2/0 to confirm they coded for the binding site of 1D10. The expression vectors used in the transfection are similar to the plasmids pVk.G and pVg.D described by Co et al. (see Co et al., *J. Immunol.* 148, 1149 (1992)), except that genes for the constant regions were derived from mouse sequences. Antibody isolated from the transfected cells was found by flow

cytometry to bind to Raji cells in a pattern indistinguishable from that of the parent mouse IgG1/kappa 1D10 antibody.

The V domains of M291 were cloned similarly and they were expressed as mouse F(ab'-zipper)<sub>2</sub> (see Kostelny et al., *J. Immunol.* 148, 1547 (1992)). Flow cytometry assay indicated M291-Fos F(ab'zipper)<sub>2</sub> binds to human T cells with similar or identical affinity as the parent antibody. This observation confirmed that the correct V domains of M291 were cloned.

## (2) Modelling and design of humanized sequences

The sequences of human V domains most similar to murine 1D10 and M291 were selected to serve as the framework of the humanized antibody. For 1D10, the best human V<sub>k</sub> sequence was R3.5H5G of human subgroup I with only sixteen differences from 1D10 in framework regions. Manheimer-Lory et al., *J. Exp. Med.* 174, 1639-1652 (1991). The best V<sub>H</sub> sequence was IC4 of Kabat's subgroup II or subgroup IV (see Kabat et al., *Sequences of Proteins of Immunological Interest* 1, 1137 (1991)), with twenty-six differences. For M291, the best human V<sub>k</sub> sequence is HF2-1/17 of human subgroup I with twenty-six amino acid differences from M291 in framework regions (Athison et al., *J. Clin. Invest.* 75, 1138 (1985); Lampman *Blood* 74, 262 (1989)); the best human V<sub>H</sub> sequence is 21/28 of human subgroup I with twenty amino acid differences. Dersimonian et al., *J. Immunol.* 139, 2496-2501 (1987). With the help of the 3-dimensional model, an additional number of framework positions that differed between the murine antibodies and the chosen human sequences were identified. The location of those amino acid residues in 3-dimensional space relative to the hypervariable regions, or CDRs, indicated they were likely to influence CDR conformation, and thus binding affinity. Murine sequences were used in these positions. A number of positions were identified in the human sequences that differed from the consensus of their respective subgroups. These amino acids were changed to correspond to consensus sequences. V<sub>H</sub> and V<sub>L</sub> sequence comparisons between the murine and humanized 1D10, and between the murine and

humanized M291, are shown in Figure 4 and Figure 5, respectively.

### (3) Synthesis and expression of humanized 1D10 antibody

DNA segments encoding the humanized 1D10 L and H chain V regions were constructed by total gene synthesis from overlapping oligonucleotides. These mini exons included signal sequences, J segments and splice donor sequences and were surrounded by XbaI sites. The DNA segments were incorporated in an expression vector using the scheme outlined in Figure 6.

The humanized V domains were cloned into the XbaI sites of the corresponding heavy and light chain expression plasmids pV<sub>G</sub>1.D.Tt and pV<sub>k</sub>.rG.dE. The resulting plasmids are called pHu1D10.Vgl.D.Tt and pHu1D10.V<sub>k</sub>.rG.dE. The heavy chain expression vector, pVgl.D.Tt, which contains the mutant dihydrofolate reductase gene (*mdhfr*) as the selectable marker (see Simonsen & Levinson, *Proc. Natl. Acad. Sci. USA*, 80, 2495, (1983)), the human cytomegalovirus (hCMV) major immediate early promoter and enhancer for transcription initiation (see Boshart et al., *Cell* 41, 521 (1985)), and the human IgG1 constant regions was constructed from the respective fragments by standard methods. It differs from the vector pV<sub>G</sub>1.D described by Co et al, *J. Immunol.* 148, 1149 (1992) by having a transcription termination site 3' to the  $\gamma$  1 gene poly(A) site. The transcription termination site (Tt) was derived from the sequence located downstream from the human complement gene C2 (+37 to +162 bp from the C2 poly(A) site) (see Ashfield et al., *EMBO J.* 10, 4197 (1991)) and was synthesized entirely by using overlapping oligonucleotides.

For light chain expression, a vector was constructed from the hCMV promoter and enhancer, the human C<sub>k</sub> gene including part of the preceding intron, and the xanthine-guanine phosphoribosyltransferase (*gpt*) gene (see Mulligan & Berg, *Proc. Natl. Acad. Sci. USA*, 78, 2072 (1981)) for selection. The vector, pV<sub>k</sub>.rG.dE, is similar to pV<sub>k</sub> described by Co et al. (see Co et al., *J. Immunol.* 148, 1149 (1992)) except for the orientation of the *gpt* gene. In



addition, one of the two repeated sequences in the enhancer region of the SV40 promoter used to transcribe the *gpt* gene was deleted by *Sph*I digestion.

For coexpression of heavy and light chains in one plasmid, an *Eco*RI fragment containing the hCMV promoter, the VH exon, the C<sub>H</sub>1, C<sub>H</sub>2 and C<sub>H</sub>3 exons, the polyA signal, and the transcription termination signal was taken from the heavy chain expression vector and cloned into the unique *Eco*RI site of the corresponding light chain expression plasmid. Due to the presence of the transcription termination signal situated between them, the two genes are transcribed independently by the hCMV promoter. After transcription the humanized V<sub>H</sub> exon is spliced to the human  $\gamma$  1 C<sub>H</sub>1, hinge, C<sub>H</sub>2 and C<sub>H</sub>3 exons, and then polyadenylated. Similarly the V<sub>L</sub> exon is spliced to the human C<sub>K</sub> exon. The predicted amino acid sequences of the mature light and heavy chains of humanized 1D10 are shown in Figures 4C and 4E, respectively.

Plasmid pHu1D10.IgG1.rG.dE, was used for transfection into mouse myeloma cell line TSO by electroporation. TSO cells are derivative of mouse myeloma NSO cells (ECACC 85110503) selected for their ability to grow in serum-free media according to the procedure of Sato et al., *J. Exp. Med.* 165, 1761 (1987). The cells from each transfection were selected for *gpt* expression. Because the SV40 promoter/enhancer for the *gpt* gene has been crippled, only few transfectants can express *gpt* high enough to survive the selection (see Jasin & Berg, *Genes Dev.* 2, 1353 (1988)). Transfection efficiency is about  $0.5-1.0 \times 10^{-6}$ ; compared to the efficiency of  $10-50 \times 10^{-6}$  from transfection using near identical plasmid containing the wild type SV40 promoter for *gpt*. When screened for production of humanized antibodies by standard ELISA, the average surviving cells also gave higher levels of antibody compared to those transfected with plasmid containing the wild type SV40 promoter. The best antibody producer was then subcloned for the production of the humanized 1D10. The antibody, Hu1D10, was purified from the serum-free spent medium by Protein A affinity chromatography.

(4) Properties of Hu1D10

Murine 1D10-IgG2a and humanized 1D10 had identical spectrums of reactivity with 1D10 positive and 1D10 negative cell lines. The affinity of murine 1D10-IgG2a and humanized 1D10 for cells bearing the target antigen was evaluated using a displacement assay (see Woodle et al., *J. Immunol.* 148 2756 (1992)). In this assay, the ability of prebound humanized 1D10 or murine 1D10-IgG2a to inhibit the binding of FITC-labeled murine 1D10-IgG2a was quantitated by FACS analysis. Humanized 1D10 competitively inhibited the binding of murine 1D10-IgG2a to a degree similar to that seen with the parent antibody (Figure 7A). These data indicated that the humanized antibody binds with similar affinity as the murine antibody. Scatchard analysis was used to better estimate the apparent affinity of humanized 1D10. Humanized 1D10-IgG1 was found to have an apparent  $K_a$  of  $2.3 \times 10^8 M^{-1}$ , and there are about  $5 \times 10^5$  sites per cell in the Raji cell line (Figure 7B). In addition, humanized 1D10 has the ability to direct ADCC and complement mediated lysis, two effector functions that are not present in the original murine 1D10 (Figures 8A and 8B).

(5) Synthesis and expression of humanized M291 and 1D10(ab'-zipper)<sub>2</sub>

Leucine zipper genes, Jun and Fos, were synthesized as described by Kostelny et al., *J. Immunol.* 148, 1547 (1992). The resulting PCR products were 179 bp PstI-SalI fragments, encompassing the entire hinge zipper gene fusion. The PstI site is the natural restriction site located at the beginning of the hinge exon, but the SalI site was added to the end of the zipper sequences during PCR. The hinge/zipper exon was inserted with a 162 bp SalI-BamHI fragment containing the 3' noncoding sequence of the mouse IgG2a gene into the heavy chain expression vector pVgl.D.Tt, replacing the hinge, C<sub>H</sub>2 and C<sub>H</sub>3 exons in the plasmid. Coexpression of the truncated heavy chain (Fd) gene with light chain gene in one plasmid is essentially the same as described above for pHu1D10.IgG1.rG.dE (Figure 6). The expression plasmids are called pHu1D10-

Jun.rG.dE and pHuM291-Fos.rG.dE (Figure 9). The differences between these plasmids and those used to express the whole antibody are: (1) the human  $\gamma 1$  C<sub>H</sub>1 exon is now spliced to the hinge/zipper fusion exon instead of the hinge, C<sub>H</sub>2 and C<sub>H</sub>3 exons, and (2) the transcript is polyadenylated by a heterologous signal. The leucine zipper Jun is used for the Fd of Hu1D10, and Fos for Fd of HuM291. When combined with the corresponding light chain, the Fd-zipper would form F(ab'-zipper)<sub>2</sub>. The humanized F(ab'-zipper)<sub>2</sub> fragments for 1D10 and M291 are called Hu1D10-Jun and HuM291-Fos, respectively. The predicted amino acid sequences of the heavy chain Fd-zipper in Hu1D10-Jun and HuM291-Fos are shown in Figures 4D and 5D, respectively. In both cases there were modifications of the human IgG1 hinge at the region of hinge/zipper fusion (Figure 10). An insertion of two amino acid residues Lys-Cys derived from the mouse IgG2a hinge was introduced to the hinge exon to provide an additional inter-heavy chain disulfide bond. The insertion of these two residues in the human IgG1 hinge renders its COOH-terminal half homologous to that of the mouse IgG2a hinge. The modified hinge would have three inter-heavy chain disulfide bonds compared to two in the wild type human IgG1. In addition an Ala residue (first residue of the C<sub>H</sub>2 domain) and two Gly residues were introduced at the fusion junction to make the joints more flexible. The expression plasmids, pHu1D10-Jun.rG.dE and pHuM291-Fos.rG.dE, were separately transfected into mouse myeloma cell line TSO by electroporation. Transfectants were screened for the presence and the quantity of secreted F(ab'-zipper)<sub>2</sub> fragments by ELISA. F(ab'-zipper)<sub>2</sub> fragments were purified using Protein G affinity chromatography.

#### (5) Properties of HuM291-Fos

The relative affinity of murine M291 and HuM291-Fos F(ab'-zipper)<sub>2</sub> for T cells was evaluated using the displacement assay described above. HuM291-Fos blocks the binding of FITC-labeled murine M291 IgG2a as well as the unlabeled M291 (Figure 11A). The affinity of HuM291 for CD3 is estimated to be within 2-3 fold of M291's. Scatchard

analysis indicated the apparent affinity of HuM291-Fos was  $K_a \sim 1.1 \times 10^9 M^{-1}$ , and there are about  $6.6 \times 10^4$  sites per cell in activated human T cells (Figure 11B).

5        (6) Formation of the bispecific Hu1D10-Jun x HuM291-Fos F(ab'-zipper)<sub>2</sub> in vitro

Hu1D10-Jun and HuM291-Fos were mixed in equal molar at concentrations between 0.5 to 3.0 mg/ml and reduced with 10 mM DTT in PBS at 37°C for 1 hour to form Fab'-zippers.

10 They were passed through Sepharose G-50 column in PBS to remove DTT. The desalted protein was incubated at 4°C for 48 hours to allow formation of heterodimeric bispecific Hu1D10-Jun x HuM291-Fos. The bispecific molecules were further purified by hydrophobic interaction chromatography (HIC) on a Phenyl Sepharose column.

15        Example 5: T cell-mediated cytotoxicity by humanized bispecific antibodies

20        The ability of Hu1D10-Jun X HuM291-Fos to direct T cell-mediated lysis was tested in a chromium-release assay. Human T cells derived from PBMC after OKT3 and IL-2 treatment were used as effector cells. Dawo, which is a cell line developed from a patient with large B cell lymphoma, was used as target cells. Figure 12 shows that the bispecific Hu1D10-Jun x HuM291-Fos, as well as the mouse bispecific IgG 1DT3-D directed T cells to lyse target cells. The two bispecific molecules seemed to have similar activities at low antibody concentrations. The two parent antibodies, HuM291-Fos and Hu1D10-Jun, were not effective in this assay, either singly or

30 in combination.

At high concentrations (10 µg/ml, 1DT3-D had higher activity than Hu1D10-Jun x HuM291-Fos in mediating target cell lysis. This was because of low affinity Fc receptors on the surface of the target cells. At high antibody concentration, bispecific IgG's Fc could bind to these receptors and direct T cell to lyse target cells independent of the target antigen, a mechanism known as reverse lysis (see Weiner et al., *J. Immunol.* 152, 2385 (1994)). Because Hu1D10-Jun x HuM291-Fos

is an  $F(ab')_2$ -like molecule without an Fc, it cannot initiate lysis by binding to an Fc receptor. In some therapeutic application, the property of the humanized antibody is advantageous in increasing selective toxicity of the antibody.

5 All publications and patent applications cited above are herein incorporated by reference in their entirety for all purposes to the same extent as if each individual publication or patent application were specifically and individually indicated to be so incorporated by reference. Although the  
10 present invention has been described in some detail by way of illustration and example for purposes of clarity and understanding, it will be apparent that certain changes and modifications may be practiced within the scope of the appended claims.  
15

WHAT IS CLAIMED IS:

1                   1.    A bispecific antibody that binds to:  
2                   (a) a first antigen on the surface of effector  
3 cells selected from the group consisting of T-cells and  
4 natural killer cells, and

5                   (b) a second antigen on a 28/32 kDa  
6 heterodimeric protein on the surface of malignant B cells,  
7 which second antigen specifically binds to an antibody  
8 designated 1D10,

9                   wherein the binding of the bispecific antibody  
10 to the first and second antigens results in killing of the  
11 malignant B cells.

1                   2.    The bispecific antibody of claim 1, wherein the  
2 effector cells are T cells.

1                   3.    The bispecific antibody of claim 2, wherein the  
2 bispecific antibody binds to a CD3 antigen on the T cells.

1                   4.    The bispecific antibody of claim 3, wherein the  
2 antibody is produced by the cell line ATCC HB 10993.

1                   5.    A cell line producing the bispecific antibody of  
2 claim 1.

1                   6.    The cell line of claim 5 that is a hybrid-  
2 hybridoma formed from two hybridomas,  
3                   a first hybridoma producing an antibody that  
4 binds to the first antigen, and  
5                   a second hybridoma producing an antibody that  
6 binds to the second antigen.

1                   7.    The cell line of claim 7, wherein the effector  
2 cells are T-cells and the first hybridoma produces an antibody  
3 that binds to a CD3 antigen on the T-cells.

1                   8.    A cell line designated ATCC HB 10993.

9. An antibody designated 1D10.

10. A humanized version of the antibody of claim 9.

11. A humanized antibody according to claim 10, the antibody comprising a humanized heavy chain and a humanized light chain:

(1) the humanized light chain comprising three complementarity determining regions (CDR1, CDR2 and CDR3) having amino acid sequences from the corresponding complementarity determining regions of the 1D10 immunoglobulin light chain, and a variable region framework from a human kappa light chain variable region framework sequence except in at least one position selected from a first group consisting of L48, L49, L69, and L70 wherein the amino acid position is occupied by the same amino acid present in the equivalent position of the 1D10 immunoglobulin light chain variable region framework; and

(2) the humanized heavy chain comprising three complementarity determining regions (CDR1, CDR2 and CDR3) having amino acid sequences from the corresponding complementarity determining regions of 1D10 immunoglobulin heavy chain, and a variable region framework from a human heavy chain variable region framework sequence except in at least one position selected from a second group consisting of H27, H29, H30, H37, H67, H71, H78 and H83, wherein the amino acid position is occupied by the same amino acid present in the equivalent position of the mouse 1D10 immunoglobulin heavy chain variable region framework;

wherein the immunoglobulin specifically binds to a 28/32 kDa heterodimeric protein on the surface of malignant B cells with a binding affinity having a lower limit of about  $10^7 \text{ M}^{-1}$  and an upper limit of about five-times the binding affinity of the 1D10 immunoglobulin.

13. The humanized antibody of claim 12, wherein the humanized light chain variable region framework is from the light chain variable region framework of the R3.5H5G antibody except in the at least one position from the first group and except at position L43, which is occupied the amino acid present in the equivalent position of a human kappa subgroup I consensus sequence;

the humanized heavy chain is from the heavy chain region variable framework of the IC4 antibody except in at least one position selected from the second group, and except at position H73, wherein the amino acid position is occupied by the same amino acid present in the equivalent position of a human immunoglobulin subgroup II or IV consensus sequence.

14. The humanized antibody of claim 13, wherein the humanized light chain comprises the amino acid sequence of Fig. 4A (upper) and

the humanized heavy chain comprises the amino acid sequence of Fig. 4B (upper).

15. The humanized antibody of claim 14, wherein the humanized light chain further comprises a human kappa constant region, the humanized heavy chain further comprises a human  $\gamma$ 1 constant region, and the humanized antibody effects ADCC and complement-mediated lysis of malignant B-cells when bound to a 28/32 kDa heterodimeric protein on the surface of the cells.

16. A humanized antibody, the antibody comprising a humanized heavy chain and a humanized light chain:

(1) the humanized light chain comprising three complementarity determining regions (CDR1, CDR2 and CDR3) having amino acid sequences from the corresponding complementarity determining regions of the mouse M291 immunoglobulin light chain, and a variable region framework from a human kappa light chain variable region framework sequence, and

(2) the humanized heavy chain comprising three complementarity determining regions (CDR1, CDR2 and CDR3)



12 having amino acid sequences from the corresponding  
13 complementarity determining regions of the mouse M291  
14 immunoglobulin heavy chain, and a variable region framework  
15 from a human heavy chain variable region framework sequence  
16 except in at least one position selected from a second group  
17 consisting of H30, H67, H68, H70, H72 and H74 wherein the  
18 amino acid position is occupied by the same amino acid present  
19 in the equivalent position of the mouse M291 immunoglobulin  
20 heavy chain variable region framework;

21 wherein the immunoglobulin specifically binds to a  
22 CD3 antigen on the surface of T cells with a binding affinity  
23 having a lower limit of about  $10^7$  M<sup>-1</sup> and an upper limit of  
24 about five-times the binding affinity of the M291  
25 immunoglobulin.

1 17. The humanized antibody of claim 16, wherein  
2 the humanized light chain variable region framework  
3 is from the light chain variable region framework of the HF2-  
4 1/17 antibody in subgroup I;

5 the humanized heavy chain region framework is from  
6 the heavy chain region variable framework of the 21/28  
7 antibody except in at least one position selected from the  
8 second group, and except at position 44, wherein the amino  
9 acid position is occupied by the same amino acid present in  
10 the equivalent position of a human immunoglobulin subgroup I  
11 consensus sequence.

1 18. The humanized antibody of claim 17, wherein  
2 the humanized light chain comprises the amino acid  
3 sequence of Fig. 5A (upper) and

4 the humanized heavy chain comprises the amino acid  
5 sequence of Fig. 5B (upper).

1 19. The bispecific antibody of claim 1 that is  
2 humanized.

1 20. The bispecific antibody of claim 18, wherein  
2 the first antigen is the CD3 antigen.

1           21. The bispecific antibody of claim 20,  
2 comprising:

3           a first binding fragment comprising:

4           a humanized form of the heavy chain variable  
5 region of the M291 antibody;

6           a humanized form of the light chain variable  
7 region of the M291 antibody; and

8           a second binding fragment, which is linked to the  
9 first binding fragment, comprising:

10          a humanized form of the heavy chain variable  
11 region from the 1D10 antibody;

12          a humanized form of the light chain variable  
13 region from the 1D10 antibody;

14          wherein the first binding fragment specifically  
15 binds to the CD3 antigen and the second binding fragment  
16 specifically binds to the 28/32 kDa heterodimeric antigen on  
17 the surface of the malignant B cells.

1           22. The bispecific antibody of claim 21, wherein:  
2           the humanized form of the heavy chain variable  
3 region of the M291 antibody comprises three complementarity  
4 determining regions (CDR1, CDR2 and CDR3) having amino acid  
5 sequences from the corresponding complementarity determining  
6 regions of M291 immunoglobulin heavy chain, and a variable  
7 region framework from a human heavy chain variable region  
8 framework sequence except in at least one position selected  
9 from a second group consisting of H30, H67, H68, H70, H72 and  
10 H74 wherein the amino acid position is occupied by the same  
11 amino acid present in the equivalent position of the mouse  
12 M291 immunoglobulin heavy chain variable region framework;

13          the humanized form of the light chain variable  
14 region of the M291 antibody comprises three complementarity  
15 determining regions (CDR1, CDR2 and CDR3) having amino acid  
16 sequences from the corresponding complementarity determining  
17 regions of the M291 immunoglobulin light chain, and a variable  
18 region framework from a human kappa light chain variable  
19 region framework sequence;

the humanized form of the heavy chain variable region from the 1D10 antibody comprises three complementarity determining regions (CDR1, CDR2 and CDR3) having amino acid sequences from the corresponding complementarity determining regions of 1D10 immunoglobulin heavy chain, and a variable region framework from a human heavy chain variable region framework sequence except in at least one position selected from a second group consisting of H27, H29, H30, H37, H67, H71, H78 and H83, wherein the amino acid position is occupied by the same amino acid present in the equivalent position of the mouse 1D10 immunoglobulin heavy chain variable region framework; and

the humanized form of the light chain variable region from the 1D10 antibody comprises three complementarity determining regions (CDR1, CDR2 and CDR3) having amino acid sequences from the corresponding complementarity determining regions of the 1D10 immunoglobulin light chain, and a variable region framework from a human kappa light chain variable region framework sequence except in at least one position selected from a first group consisting of L48, L49, L69, and L70 wherein the amino acid position is occupied by the same amino acid present in the equivalent position of the 1D10 immunoglobulin light chain variable region framework.

23. The bispecific antibody of claim 22, wherein the first binding fragment comprises the heavy chain variable region shown in Fig. 5B (upper) and the light chain variable region shown in Fig. 5A (upper), and the second binding fragment comprises the heavy chain variable region shown in Fig. 4B (upper) and the light chain variable region shown in Fig. 4A (upper).

24. The bispecific antibody of claim 23, wherein the first and second binding fragments each further comprises a segment of a constant region fused to the respective heavy chain variable regions, and the binding fragments are linked by association of the constant regions.

1           25. The bispecific antibody of claim 24, wherein  
2 the binding fragments are Fab or Fab'.

1           26. The bispecific antibody of claim 25, wherein  
2 the first and second binding fragments are Fab's and the  
3 bispecific antibody is a F(ab')<sub>2</sub>.

1           27. The bispecific antibody of claim 26, wherein  
2 the first and second binding fragments further comprise first  
3 and second leucine zippers fused to the respective constant  
4 regions.

1           28. The bispecific antibody of claim 27, wherein  
2 the first binding fragment comprises a heavy chain having the  
3 amino acid sequence shown in Fig. 5D and the second binding  
4 fragment comprises a heavy chain having the amino acid  
5 sequence shown in Fig. 4D.

1           29. A method of treating a patient suffering from  
2 malignant B-cells, the method comprising treating the patient  
3 with a therapeutically effective amount of the bispecific  
4 antibody of claim 1.

1           30. The method of claim 29, wherein the bispecific  
2 antibody comprises:

3           a first binding fragment comprising:

4               a humanized form of the heavy chain variable  
5 region of the M291 antibody;

6               a humanized form of the light chain variable  
7 region of the M291 antibody; and

8               a second binding fragment, which is linked to the  
9 first binding fragment, comprising:

10               a humanized form of the heavy chain variable  
11 region from the 1D10 antibody;

12               a humanized form of the light chain variable  
13 region from the 1D10 antibody;

14               wherein the first binding fragment specifically  
15 binds to the CD3 antigen on the surface of T-cells and the



## BISPECIFIC ANTIBODY EFFECTIVE TO TREAT B-CELL LYMPHOMA AND CELL LINE

### ABSTRACT OF THE DISCLOSURE

5       The invention provides bispecific antibodies with  
selective cytotoxicity against malignant B-cells. The  
bispecific antibodies bind to an effector cell antigen and to  
a 28/32 kDa heterodimeric protein on the surface of malignant  
B-cells. The invention also includes the monospecific  
10       components of the bispecific antibodies, humanized versions  
thereof, and humanized bispecific antibodies. The invention  
further provides therapeutic and diagnostic methods employing  
these antibodies.




FIG. 1




FIG. 3




FIG. 2











1 D I Q M T Q S P S S L S A S V G D R V T  
 21 I T C S A S S S V S Y M N W Y Q Q K P G  
 41 K A P K R L I Y D T S K L A S G V P S R  
 61 F S G S G S G T D F T L T I S S L Q P E  
 81 D F A T Y Y C Q Q W S S N P P T F G G G  
 101 T K V E I K R T V A A P S V F I F P P S  
 121 D E Q L K S G T A S V V C L L N N F Y P  
 141 R E A K V Q W K V D N A L Q S G N S Q E  
 161 S V T E Q D S K D S T Y S L S S T L T L  
 181 S K A D Y E K H K V Y A C E V T H Q G L  
 201 S S P V T K S F N R G E C.

FIG. 5C

1 Q V Q L V Q S G A E V K K P G A S V K V  
 21 S C K A S G Y T F I S Y T M H W V R Q A  
 41 P G Q G L E W M G Y I N P R S G Y T H Y  
 61 N Q K L K D K A T L T A D K S A S T A Y  
 81 M E L S S L R S E D T A V Y Y C A R S A  
 101 Y Y D Y D G F A Y W G Q G T L V T V S S  
 121 A S T K G P S V F P L A P S S K S T S G  
 141 G T A A L G C L V K D Y F P E P V T V S  
 161 W N S G A L T S G V H T F P A V L Q S S  
 181 G L Y S L S S V V T V P S S S L G T Q T  
 201 Y I C N V N H K P S N T K V D K K V E P  
 221 K S C D K T H T C P P C K C P A G G L T  
 241 D T L Q A E T D Q L E D K K S A L Q T E  
 261 I A N L L K E K E K L E F I L A A T S

FIG. 5D






FIG. 7A




FIG. 7B





FIG. 8A









FIG. 9

A-E-P-K-S-C-D-K-T-H-T-C-P-P-C-K-C-P-A-G-G-Zipper  
———— Modified Human IgG1 Hinge ————

FIG. 10A

A-E-P-K-S-C-D-K-T-H-T-C-P-P-C-P-  
———— Human IgG1 Hinge ————

FIG. 10B

E-P-R-G-P-T-I-K-P-C-P-P-C-K-C-P-  
———— Mouse IgG2a Hinge ————

FIG. 10C




FIG. 11A




FIG. 11B

Lysis of Dawo cells mediated by bispecific antibody




FIG. 12

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to:  
Commissioner of Patents and Trademarks,  
Attn: Application Processing Division  
Special Processing and Correspondence Branch  
Washington, D.C. 20231 on June 27, 1995

By Joe S. Winter

Attorney Docket No. 11823-004910

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:	)	Examiner: Unknown
	)	
GEORGE WEINER ET AL.	)	Art Unit: Unknown
	)	
Serial No.: 08/397,411	)	
	)	
Filed: March 1, 1995	)	
	)	
For: BISPECIFIC ANTIBODY	)	TRANSMITTAL OF SEQUENCE
EFFECTIVE TO TREAT B-CELL	)	<u>LISTING</u>
LYMPHOMA AND CELL LINE	)	
	)	

Commissioner of Patents and Trademarks  
Washington, D.C. 20231

Sir:

Applicants hereby submit a "Sequence Listing" and a copy of same in computer-readable form.

Applicants attest that the information in the "Sequence Listing" is identical to that which is in computer-readable form, as required under 37 CFR 1.821(f). This submission introduces no new matter, since the enclosed sequences are identical to the sequences which were submitted in the original patent application. Two copies of this paper are enclosed.

Respectfully submitted,

Joe Liebeschuetz

Joe Liebeschuetz  
Reg. No. 37,505

TOWNSEND and TOWNSEND and CREW  
20th Floor, Steuart Street Tower  
One Market Plaza  
San Francisco, California 94105-1492  
(415) 326-2400  
11823/49-1.TSQ



[illegible]

(i) SEQUENCE CHARACTERISTICS:

SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 107 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly  
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Glu Asn Ile Tyr Ser Tyr  
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Val  
35 40 45

Ser Asn Ala Lys Thr Leu Ala Glu Gly Val Pro Ser Arg Phe Ser Gly  
50 55 60

Ser Gly Ser Gly Lys Gln Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro  
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln His His Tyr Gly Asn Ser Tyr  
85 90 95

Pro Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys  
100 105

Variable	Mean	SD	Min	Max
Age	35.2	12.5	18	65
Gender	50.0	50.0	0	100
Marital Status	65.0	25.0	0	100
Education	12.5	2.0	8	16
Income	45000	15000	20000	80000
Health Status	75.0	15.0	50	100
Stress Level	60.0	20.0	30	90
Sleep Quality	70.0	18.0	40	100
Dietary Habits	68.0	22.0	40	100
Exercise Frequency	55.0	25.0	20	100
Work-Life Balance	62.0	20.0	30	100
Family Support	78.0	15.0	50	100
Community Involvement	65.0	20.0	30	100
Personal Growth	72.0	18.0	40	100
Life Satisfaction	75.0	15.0	50	100
Overall Well-being	70.0	18.0	40	100

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 107 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Asp 1	Ile	Gln	Met 5	Thr	Gln	Ser	Pro	Ala	Ser 10	Leu	Ser	Ala	Ser	Val 15	Gly
Glu	Thr	Val	Thr 20	Ile	Thr	Cys	Arg	Ala 25	Ser	Glu	Asn	Ile	Tyr 30	Ser	Tyr
Leu	Ala	Trp 35	Tyr	Gln	Gln	Lys	Gln 40	Gly	Lys	Ser	Pro	Gln 45	Leu	Leu	Val
Ser	Asn 50	Ala	Lys	Thr	Leu	Ala 55	Glu	Gly	Val	Thr	Ser 60	Arg	Phe	Ser	Gly
Ser 65	Gly	Ser	Gly	Lys	Gln 70	Phe	Ser	Leu	Lys	Ile 75	Asn	Ser	Leu	Gln	Pro 80
Glu	Asp	Phe	Ala	Thr 85	Tyr	Tyr	Cys	Gln	His 90	His	Tyr	Gly	Asn	Ser 95	Tyr
Pro	Phe	Gly	Gly 100	Gly	Thr	Lys	Leu	Glu 105	Ile	Lys					



[illegible]

(A) LENGTH: 116 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

[illegible]

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 116 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Gln Val Gln Leu Lys Gln Ser Gly Pro Gly Leu Val Gln Pro Ser Gln  
1 5 10 15  
Ser Leu Ser Ile Thr Cys Thr Gly Ser Gly Phe Ser Leu Thr Asn Tyr  
20 25 30  
Gly Val His Trp Val Arg Gln Ser Pro Gly Lys Gly Leu Glu Trp Ile  
35 40 45  
Gly Val Lys Trp Ser Gly Gly Ser Thr Glu Tyr Asn Ala Ala Phe Ile  
50 55 60  
Ser Arg Leu Ser Ile Ser Lys Asp Asn Ser Lys Ser Gln Val Phe Phe  
65 70 75 80  
Lys Met Asn Ser Leu Gln Ala Asp Asp Thr Ala Met Tyr Tyr Cys Ala  
85 90 95  
Arg Asn Asp Arg Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr Ser Val  
100 105 110  
Thr Val Ser Ser  
115

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 214 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly  
1 5 10 15  
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Glu Asn Ile Tyr Ser Tyr  
20 25 30  
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Val  
35 40 45  
Ser Asn Ala Lys Thr Leu Ala Glu Gly Val Pro Ser Arg Phe Ser Gly  
50 55 60  
Ser Gly Ser Gly Lys Gln Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro  
65 70 75 80  
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln His His Tyr Gly Asn Ser Tyr  
85 90 95  
Pro Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala  
100 105 110  
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly  
115 120 125  
Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala  
130 135 140  
Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln  
145 150 155 160  
Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser  
165 170 175  
Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr  
180 185 190  
Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser  
195 200 205

CCF000000000





Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr
210						215					220				
Cys	Pro	Pro	Cys	Lys	Cys	Pro	Ala	Gly	Gly	Arg	Ile	Ala	Arg	Leu	Glu
225					230					235					240
Glu	Lys	Val	Lys	Thr	Leu	Lys	Ala	Gln	Asn	Ser	Glu	Leu	Ala	Ser	Thr
				245					250					255	
Ala	Asn	Met	Leu	Arg	Glu	Gln	Val	Ala	Gln	Leu	Lys	Gln	Lys	Val	Met
			260					265					270		

Asn

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 446 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Gln	Val	Gln	Leu	Gln	Glu	Ser	Gly	Pro	Gly	Leu	Val	Lys	Pro	Ser	Glu	
1				5					10					15		
Thr	Leu	Ser	Leu	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Asn	Tyr	
			20					25					30			
Gly	Val	His	Trp	Val	Arg	Gln	Ser	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Ile	
		35					40					45				
Gly	Val	Lys	Trp	Ser	Gly	Gly	Ser	Thr	Glu	Tyr	Asn	Ala	Ala	Phe	Ile	
		50				55					60					
Ser	Arg	Leu	Thr	Ile	Ser	Lys	Asp	Thr	Ser	Lys	Asn	Gln	Val	Ser	Leu	
65					70					75					80	
Lys	Leu	Asn	Ser	Leu	Thr	Ala	Ala	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	
				85					90					95		
Arg	Asn	Asp	Arg	Tyr	Ala	Met	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	
			100					105					110			
Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	
			115				120					125				
Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	
			130			135					140					
Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	
145					150					155					160	
Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	
				165					170					175		
Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	
			180					185					190			
Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	
			195				200					205				





(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 106 amino acids  
    (B) TYPE: amino acid  
    (C) STRANDEDNESS: single  
    (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
1				5				10						15	
Asp	Arg	Val	Thr	Ile	Thr	Cys	Ser	Ala	Ser	Ser	Ser	Val	Ser	Tyr	Met
			20					25					30		
Asn	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Arg	Leu	Ile	Tyr
			35				40					45			
Asp	Thr	Ser	Lys	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser
			50			55					60				
Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Glu
65					70					75					80
Asp	Phe	Asp	Thr	Tyr	Tyr	Cys	Gln	Gln	Trp	Ser	Ser	Asn	Pro	Pro	Thr
				85					90					95	
Phe	Gly	Gly	Gly	Thr	Lys	Val	Glu	Ile	Lys						
			100					105							

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 106 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Asp	Ile	Val	Leu	Thr	Gln	Ser	Pro	Ala	Ile	Met	Ser	Ala	Ser	Pro	Gly
1				5					10					15	
Glu	Lys	Val	Thr	Met	Thr	Cys	Ser	Ala	Ser	Ser	Ser	Val	Ser	Tyr	Met
			20					25					30		
Asn	Trp	Tyr	Gln	Gln	Lys	Ser	Gly	Thr	Ser	Pro	Lys	Arg	Trp	Thr	Tyr
		35					40					45			
Asp	Thr	Ser	Lys	Leu	Ala	Ser	Gly	Val	Pro	Ala	Arg	Phe	Ser	Gly	Ser
	50					55					60				
Gly	Ser	Gly	Thr	Ser	Tyr	Ser	Leu	Thr	Ile	Ser	Ser	Met	Glu	Ala	Glu
65					70					75					80
Asp	Ala	Asp	Thr	Tyr	Tyr	Cys	Gln	Gln	Trp	Ser	Ser	Asn	Pro	Pro	Thr
			85						90				95		
Phe	Gly	Ser	Gly	Thr	Lys	Leu	Glu	Ile	Lys						
			100					105							



Variable	Unit	Value
Age	Years	65.0
Sex	Male/Female	50/50
Weight	kg	70.0
Height	m	1.75
BMI	kg/m²	22.2
Heart rate	beats/min	72.0
Blood pressure	mmHg	120/80
Cholesterol	mmol/L	5.2
Glucose	mmol/L	5.0
Hemoglobin	g/dL	15.0
Hematocrit	%	45.0
White blood cells	10³/mm³	7.0
Platelets	10³/mm³	250.0
Neutrophils	%	60.0
Lymphocytes	%	35.0
Monocytes	%	5.0
Eosinophils	%	0.5
Basophils	%	0.5
Prothrombin time	sec	12.0
Partial thromboplastin time	sec	30.0
Fibrinogen	g/L	3.0
D-dimer	ng/mL	0.5
C-reactive protein	mg/L	1.0
Erythrocyte sedimentation rate	mm/hr	10.0
Urea nitrogen	mg/dL	10.0
Creatinine	mg/dL	1.0
BUN/Cr ratio	mg/dL	10.0
Calcium	mg/dL	10.0
Phosphorus	mg/dL	3.0
Magnesium	mg/dL	2.0
Sodium	mg/dL	140.0
Potassium	mg/dL	4.0
Chloride	mg/dL	100.0
Bicarbonate	mg/dL	24.0
Lactate	mg/dL	1.0
Ammonia	mg/dL	10.0
Bilirubin	mg/dL	1.0
Aspartate aminotransferase	U/L	40.0
Alanine aminotransferase	U/L	40.0
Gamma-glutamyl transferase	U/L	40.0
Alkaline phosphatase	U/L	40.0
Prothrombin time	sec	12.0
Partial thromboplastin time	sec	30.0
Fibrinogen	g/L	3.0
D-dimer	ng/mL	0.5
C-reactive protein	mg/L	1.0
Erythrocyte sedimentation rate	mm/hr	10.0
Urea nitrogen	mg/dL	10.0
Creatinine	mg/dL	1.0
BUN/Cr ratio	mg/dL	10.0
Calcium	mg/dL	10.0
Phosphorus	mg/dL	3.0
Magnesium	mg/dL	2.0
Sodium	mg/dL	140.0
Potassium	mg/dL	4.0
Chloride	mg/dL	100.0
Bicarbonate	mg/dL	24.0
Lactate	mg/dL	1.0
Ammonia	mg/dL	10.0
Bilirubin	mg/dL	1.0
Aspartate aminotransferase	U/L	40.0
Alanine aminotransferase	U/L	40.0
Gamma-glutamyl transferase	U/L	40.0
Alkaline phosphatase	U/L	40.0

(A) LENGTH: 120 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Gln 1	Val	Gln	Leu 5	Gln	Gln	Ser	Gly	Ala	Glu 10	Leu	Ala	Arg	Pro	Gly 15	Ala
Ser	Val	Lys	Met 20	Ser	Cys	Lys	Ala	Ser 25	Gly	Tyr	Thr	Phe	Ile 30	Ser	Tyr
Thr	Met	His 35	Trp	Val	Lys	Gln	Arg 40	Pro	Gly	Gln	Gly	Leu 45	Glu	Trp	Ile
Gly	Tyr 50	Ile	Asn	Pro	Arg	Ser 55	Gly	Tyr	Thr	His	Tyr 60	Asn	Gln	Lys	Leu
Lys 65	Asp	Lys	Ala	Thr	Leu 70	Thr	Ala	Asp	Lys	Ser 75	Ser	Ser	Ser	Ala	Tyr 80
Met	Gln	Leu	Ser	Ser 85	Leu	Thr	Ser	Glu	Asp 90	Ser	Ala	Val	Tyr	Tyr 95	Cys
Ala	Arg	Ser	Ala 100	Tyr	Tyr	Asp	Tyr	Asp 105	Gly	Phe	Ala	Tyr	Trp 110	Gly	Gln
Gly	Thr	Leu 115	Val	Thr	Val	Ser	Ala 120								

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 213 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly	
1				5				10						15		
Asp	Arg	Val	Thr	Ile	Thr	Cys	Ser	Ala	Ser	Ser	Ser	Val	Ser	Tyr	Met	
			20					25					30			
Asn	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Arg	Leu	Ile	Tyr	
		35					40					45				
Asp	Thr	Ser	Lys	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	
		50				55					60					
Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Glu	
65					70					75					80	
Asp	Phe	Asp	Thr	Tyr	Tyr	Cys	Gln	Gln	Trp	Ser	Ser	Asn	Pro	Pro	Thr	
				85					90				95			
Phe	Gly	Gly	Gly	Thr	Lys	Val	Glu	Ile	Lys	Arg	Thr	Val	Ala	Ala	Pro	
			100					105					110			
Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	
		115					120					125				
Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	
		130				135					140					
Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	
145					150					155					160	
Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	
				165					170					175		
Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	
			180					185					190			
Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	
		195					200					205				

Asn Arg Gly Glu Cys  
210

[illegible]

(A) LENGTH: 279 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Gln 1	Val	Gln	Leu	Val 5	Gln	Ser	Gly	Ala	Glu 10	Val	Lys	Lys	Pro	Gly 15	Ala
Ser	Val	Lys	Val 20	Ser	Cys	Lys	Ala	Ser 25	Gly	Tyr	Thr	Phe	Ile 30	Ser	Tyr
Thr	Met	His 35	Trp	Val	Arg	Gln	Ala 40	Pro	Gly	Gln	Gly	Leu 45	Glu	Trp	Met
Gly	Tyr 50	Ile	Asn	Pro	Arg	Ser 55	Gly	Tyr	Thr	His	Tyr 60	Asn	Gln	Lys	Leu
Lys 65	Asp	Lys	Ala	Thr	Leu 70	Thr	Ala	Asp	Lys	Ser 75	Ala	Ser	Thr	Ala	Tyr 80
Met	Glu	Leu	Ser	Ser 85	Leu	Arg	Ser	Glu	Asp 90	Thr	Ala	Val	Tyr	Tyr 95	Cys
Ala	Arg	Ser	Ala 100	Tyr	Tyr	Asp	Tyr	Asp 105	Gly	Phe	Ala	Tyr	Trp 110	Gly	Gln
Gly	Thr	Leu 115	Val	Thr	Val	Ser	Ser 120	Ala	Ser	Thr	Lys	Gly 125	Pro	Ser	Val
Phe	Pro 130	Leu	Ala	Pro	Ser	Ser 135	Lys	Ser	Thr	Ser	Gly 140	Gly	Thr	Ala	Ala
Leu 145	Gly	Cys	Leu	Val	Lys 150	Asp	Tyr	Phe	Pro	Glu 155	Pro	Val	Thr	Val	Ser 160
Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val
				165					170					175	
Leu	Gln	Ser	Ser 180	Gly	Leu	Tyr	Ser	Leu 185	Ser	Ser	Val	Val	Thr 190	Val	Pro
Ser	Ser	Ser 195	Leu	Gly	Thr	Gln	Thr 200	Tyr	Ile	Cys	Asn	Val 205	Asn	His	Lys





[illegible]

(ii) MOLECULE TYPE: peptide

Leu Xaa Xaa Xaa Xaa Xaa Xaa  
1 5